ON THE BRAIDED CONNES-MOSCOVICI CONSTRUCTION

IVAN BARTULOVIĆ

ABSTRACT. In 1998, Connes and Moscovici defined the cyclic cohomology of Hopf algebras. In 2010, Khalkhali and Pourkia proposed a braided generalization: to any Hopf algebra H in a braided category \mathcal{B} , they associate a paracocyclic object in \mathcal{B} . In this paper we explicitly compute the powers of the paracocyclic operator of this paracocyclic object. Also, we introduce twisted modular pairs in involution for H and derive (co)cyclic modules from them. Finally, we relate the paracocyclic object associated with H to that associated with an H-module coalgebra via a categorical version of the Connes-Moscovici trace.

CONTENTS

1

т.	minoduction	1
2.	Preliminaries on monoidal categories and braided Hopf algebras	3
3.	Simplicial, paracyclic, and cyclic objects	11
4.	Modular pairs and braided Connes-Moscovici construction	13
5.	Categorical Connes-Moscovici trace	17
6.	Proof of Theorem 2	23
7.	Proof of Theorem 7	35
8.	Appendix	38
References		41

1. INTRODUCTION

Cyclic (co)homology of algebras was introduced in the 1980s by Connes [3, 4] and Tsygan [11] independently. To any algebra is associated a cocyclic vector space (that is, a cocyclic object in the category of vector spaces) whose cohomology is called the cyclic cohomology of the algebra. The notion of a (co)cyclic object in a category, introduced by Connes [3], is a generalization of the notion of a (co)simplicial object in that category.

Cyclic cohomology has been considered in various versions and generalizations. In particular, in [6], Connes and Moscovici defined the Hopf cyclic cohomology by associating a cocyclic vector space to a Hopf algebra H over \mathbb{C} endowed with a modular pair in involution (that is, a pair (δ, σ) where $\delta \colon H \to \mathbb{C}$ is a character and $\sigma \in H$ is a grouplike element verifying the modular pair condition $\delta(\sigma) = 1$ and a certain involutivity condition). Also, in [5], they relate the Hopf cyclic cohomology of H to the cyclic cohomology of an H-module algebra by means of a trace map.

Braided monoidal categories were defined by Street and Joyal in the 1980s and appeared in many areas of mathematics such as low-dimensional topology and representation theory. Several generalizations of cyclic (co)homology were introduced in the braided

Introduction

²⁰²⁰ Mathematics Subject Classification. 16T05, 18M15.

Keywords. Hopf algebras, braided monoidal categories, traces.

IVAN BARTULOVIĆ

setting. In this paper, we focus on the braided generalization of the Connes-Moscovici construction due to Khalkhali and Pourkia [8]. Let H be a Hopf algebra in a braided monoidal category $\mathcal{B} = (\mathcal{B}, \otimes, \mathbb{1})$. A modular pair for H is a pair (δ, σ) , where $\delta \colon H \to \mathbb{1}$ is an algebra morphism and $\sigma \colon \mathbb{1} \to H$ is a coalgebra morphism such that $\delta \sigma = \mathrm{id}_{\mathbb{1}}$. In [8], Khalkhali and Pourkia associate to any modular pair (δ, σ) for H a paracocyclic object $\mathbf{CM}_{\bullet}(H, \delta, \sigma) = {\mathbf{CM}_n(H, \delta, \sigma)}_{n \in \mathbb{N}}$ in \mathcal{B} . This object is cocyclic if it satisfies the cocyclicity condition: for all $n \in \mathbb{N}$,

$$(\tau_n(\delta,\sigma))^{n+1} = \mathrm{id}_{H^{\otimes n}},$$

where $\tau_{\bullet}(\delta, \sigma) = {\tau_n(\delta, \sigma)}_{n \in \mathbb{N}}$ is the paracocyclic operator of $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$. As already noticed in [6] (corresponding to the case $\mathcal{B} = \operatorname{Vect}_{\mathbb{C}}$), verifying the cocyclicity condition (if true) is a rather technical task. Khalkhali and Pourkia proved [8, Theorem 7.3] that if (δ, σ) is a so called braided modular pair in involution, then $(\tau_2(\delta, \sigma))^3$ is equal to the square of the braiding of H with itself. In particular, if \mathcal{B} is symmetric, then $(\tau_2(\delta, \sigma))^3 = \operatorname{id}_{H^{\otimes 2}}$. They also state a similar claim about $(\tau_n(\delta, \sigma))^{n+1}$, which implies the cocyclicity condition when \mathcal{B} is symmetric, see [8, Remark 7.4].

Our first main result is a complete computation (by means of the Penrose graphical calculus) of the powers (up to n+1) of the paracocyclic operator $\tau_n(\delta, \sigma)$ associated with a modular pair (δ, σ) for H, see Theorem 2. Next, assume that \mathcal{B} has a twist θ . We introduce the notion of a θ -twisted modular pair in involution for H (see Section 4.1) and prove (see Corollary 4) that if (δ, σ) is such a pair, then the associated paracocyclic operator satisfies the following twisted cocyclicity condition: for all $n \in \mathbb{N}$,

$$(\tau_n(\delta,\sigma))^{n+1} = \theta_{H^{\otimes n}}.$$

When \mathcal{B} is further k-linear, we derive (co)cyclic k-modules from a θ -twisted modular pair (δ, σ) by composing $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ with the functors $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, -)$ and $\operatorname{Hom}_{\mathcal{B}}(-, \mathbb{1})$, see Section 4.3. Note that if \mathcal{B} is symmetric, then a braided modular pair in involution in the sense of [8] is a id_B-twisted modular pair in involution, where id_B is the trivial twist of \mathcal{B} , and so the associated paracocyclic operator satisfies the cocyclicity condition.

Let H be a Hopf algebra in braided category \mathcal{B} with a twist θ . Our second main result is the construction of traces à la Connes-Moscovici. More precisely, let C be a H-module coalgebra, that is, a coalgebra in the category of right H-modules in \mathcal{B} . Inspired by a construction of Akrami and Majid [1], we associate to C a paracocyclic object $\mathbf{C}_{\bullet}(C)$ in \mathcal{B} . We introduce the notion of a δ -invariant σ -trace for C and derive from each such trace a natural transformation from $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ to $\mathbf{C}_{\bullet}(C)$, see Theorem 7. This generalizes the standard Connes-Moscovici trace. We provide examples of traces in the case where \mathcal{B} is a ribbon category and H is its coend (see Section 5.3).

The paper is organized as follows. In Section 2, we review braided monoidal categories, Hopf algebras, and graphical calculus. Section 3 is devoted to preliminaries on simplicial, paracyclic, and cyclic objects in a category. In Sections 4 and 5, we state our main results and their corollaries. Sections 6 and 7 are devoted to the proofs of Theorems 2 and 7. In Appendix, we provide an alternative proof (by using the Penrose graphical calculus) of the fact that the object $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ defined in [8] is paracocyclic.

Throughout the paper, \Bbbk denotes any commutative ring. The class of objects of a category \mathcal{B} is denoted by $Ob(\mathcal{B})$.

Acknowledgments. This work was supported by the Labex CEMPI (ANR-11-LABX-0007-01), by the Région Hauts-de-France, and by the FNS-ANR grant OChoTop (ANR-18-CE93-0002-01). The author is particularly thankful for the useful advices of his PhD advisor Alexis Virelizier.

2. Preliminaries on monoidal categories and braided Hopf algebras

In this section, we recall some algebraic preliminaries used in the paper. We first recall some facts about braided monoidal categories and the Penrose graphical calculus. Next, we recall definitions of categorical Hopf algebras and related concepts. We finish with a recall on pivotal categories and coends. For a more comprehensive introduction, see [12].

2.1. Conventions. In what follows, we suppress in our formulas the associativity and unitality constraints of the monoidal category. This does not lead to any ambiguity since Mac Lane's coherence theorem (see [9]) implies that all possible ways of inserting these constraints give the same results. We will denote by \otimes and $\mathbb{1}$ the monoidal product and unit object of a monoidal category. For any objects X_1, \ldots, X_n of a monoidal category with $n \geq 2$, we set

$$X_1 \otimes X_2 \otimes \cdots \otimes X_n = (\cdots ((X_1 \otimes X_2) \otimes X_3) \otimes \cdots \otimes X_{n-1}) \otimes X_n$$

and similarly for morphisms. A monoidal category is k-linear if its Hom sets have a structure of a k-module such that the composition and monoidal product of morphisms are k-bilinear.

2.2. Braided categories. A braiding of a monoidal category $(\mathcal{B}, \otimes, \mathbb{1})$ is a family $\tau = \{\tau_{X,Y} \colon X \otimes Y \to Y \otimes X\}_{X,Y \in Ob(\mathcal{B})}$ of natural isomorphisms such that

(1)
$$\tau_{X,Y\otimes Z} = (\mathrm{id}_Y \otimes \tau_{X,Z})(\tau_{X,Y} \otimes \mathrm{id}_Z) \text{ and}$$

(2) $\tau_{X\otimes Y,Z} = (\tau_{X,Z} \otimes \mathrm{id}_Y)(\mathrm{id}_X \otimes \tau_{Y,Z})$

for all $X, Y, Z \in Ob(\mathcal{B})$. A braided category is a monoidal category endowed with a braiding. A braiding τ of \mathcal{B} is symmetric if for all $X, Y \in Ob(\mathcal{B})$,

$$\tau_{Y,X}\tau_{X,Y}=\mathrm{id}_{X\otimes Y}.$$

A symmetric category is a category endowed with a symmetric braiding.

2.3. Twists for braided categories. A *twist* for a braided monoidal category \mathcal{B} is a natural isomorphism $\theta = \{\theta_X : X \to X\}_{X \in Ob(\mathcal{B})}$ such that

(3)
$$\theta_{X\otimes Y} = (\theta_X \otimes \theta_Y) \tau_{Y,X} \tau_{X,Y}$$

holds for all $X, Y \in Ob(\mathcal{B})$. Note that this implies that $\theta_1 = id_1$. For example, any ribbon category (see Section 2.16) has a canonical twist.

Note that when \mathcal{B} is symmetric, a twist for \mathcal{B} is nothing but a monoidal natural isomorphism of the identity functor $\mathcal{B} \to \mathcal{B}$. In particular, $\mathrm{id}_{\mathcal{B}} = {\mathrm{id}_X : X \to X}_{X \in \mathrm{Ob}(\mathcal{B})}$ is a twist for \mathcal{B} .

By a *braided category with a twist*, we mean a braided category endowed with a twist.

IVAN BARTULOVIĆ

2.4. Graphical calculus. Throughout this paper, we will use the *Penrose graphical calculus*. For a systematic treatment, one may consult [12]. The diagrams are to be read from bottom to top. In a monoidal category \mathcal{B} , the diagrams are made of arcs colored by objects of \mathcal{B} and of boxes, colored by morphisms of \mathcal{B} . Arcs colored by 1 may be omitted in the pictures. The identity morphism of an $\overrightarrow{object X}$, a morphism $f: X \to Y$ in \mathcal{B} , and its composition with a morphism $g: Y \to Z$ in \mathcal{B} are represented respectively as



The tensor product of two morphisms $f: X \to Y$ and $g: U \to V$ is represented by placing a picture of f to the left of the picture of g: PSfrag replacements

$$f \otimes g = \begin{bmatrix} Y & V \\ f & g \\ X & U \end{bmatrix}$$

Any diagram represents a morphism. For example, the diagram



represents $(g \otimes \mathrm{id}_V)(f \otimes \mathrm{id}_T \otimes h)(\mathrm{id}_X \otimes c) \colon X \to Z \otimes V$. The morphism associated to a diagram depends only on the isotopy class of the diagram representing it. For example, the following *level-exchange property*: $f \qquad f \qquad g \qquad g$

reflects the formula

$$f \otimes g = (f \otimes \mathrm{id}_V)(\mathrm{id}_X \otimes g) = (\mathrm{id}_Y \otimes g)(f \otimes \mathrm{id}_U).$$

When \mathcal{B} is braided with braiding τ , we depict

$$\underbrace{ \underset{X \in Y}{\text{PSfrag replacements}}}_{Y \in X} \underbrace{ \begin{array}{c} Y \in X \\ \text{PSfrag replacements} \\ \text{and} \end{array}}_{X \in Y} \underbrace{ \tau_{X,Y}^{-1} = \stackrel{X}{Y}}_{X \in Y} \underbrace{ \begin{array}{c} Y \in X \\ \text{A} \\ Y \end{array}}_{X \in Y} .$$



Naturality of the braiding and the level-exchange property imply that for any two morphisms $f: X \to Y$ and $g: U \to V_U$ in \mathcal{B} , $U \to U$



When \mathcal{B} is braided with a twist $\theta = \{\theta_X \colon X \to X\}_{X \in Ob(\mathcal{B})}$, we denote the twist by

Axiom (3) for θ gives that for any $X, Y \in Ob(\mathcal{B})$,

$$\underbrace{\frac{\mathrm{PSfrag}}{\mathrm{PSfrag}}}_{\mathrm{PSfrag replacements}} X \otimes Y \bigcup_{X \otimes Y} X \otimes Y \bigcup_{X \in Y} X = X \otimes Y$$

2.5. Categorical algebras. An algebra in a monoidal category \mathcal{B} is a triple (A, m, u), where A is an object of \mathcal{B} , $m: A \otimes A \to A$ and $u: \mathbb{1} \to A$ are morphisms in \mathcal{B} , called *multiplication* and *unit* respectively, which satisfy the associativity and unitality axioms:

$$m(m \otimes \operatorname{id}_A) = m(\operatorname{id}_A \otimes m) \quad ext{and} \quad m(u \otimes \operatorname{id}_A) = \operatorname{id}_A = m(\operatorname{id}_A \otimes u).$$

The multiplication and unit are depicted by

$$m = \bigcup_{i=1}^{n} \text{ and } u = \bigcup_{i=1}^{n},$$

so that the associativity and unitality axioms rewrite graphically as

Here, it is understood that the arcs are colored by the underlying object of the algebra.

An algebra morphism between algebras (A, m, u) and (A', m', u') in a monoidal category \mathcal{B} is a morphism $f: A \to A'$ in \mathcal{B} such that $fm = m'(f \otimes f)$ and fu = u'. The latter conditions are depicted by

IVAN BARTULOVIĆ

2.6. Categorical coalgebras. A *coalgebra* in a monoidal category \mathcal{B} is given by a triple (C, Δ, ε) , where C is an object of \mathcal{B} , $\Delta: C \to C \otimes C$ and $\varepsilon: C \to \mathbb{1}$ are morphisms in \mathcal{B} , called *comultiplication* and *counit* respectively, which satisfy the coassociativity and counitality axioms:

 $(\Delta \otimes \mathrm{id}_C)\Delta = (\mathrm{id}_C \otimes \Delta)\Delta$ and $(\mathrm{id}_C \otimes \varepsilon)\Delta = \mathrm{id}_C = (\varepsilon \otimes \mathrm{id}_C)\Delta.$

The comultiplication and counit are depicted by

$$\underline{\frac{PSfrag replacements}{C}}_{C} \xrightarrow{PSfrag replacements}}_{C} ,$$

so that the coassociativity and counitality axioms rewrite graphically as

$$\begin{array}{c} & & \\ & &$$

A coalgebra morphism between coalgebras (C, Δ, ε) and $(C', \Delta', \varepsilon')$ in a monoidal category \mathcal{B} is a morphism $f: C \to C'$ in \mathcal{B} such that $\Delta' f = (f \otimes f) \Delta$ and $\varepsilon f = \varepsilon'$.

2.7. Graphical calculus and iterated (co)multiplications. Let (A, m, u) and (C, Δ, ε) be an algebra and a coalgebra in a monoidal category \mathcal{B} . For any $n \in \mathbb{N}$, we define the *n*-th multiplication $m_n: A^{\otimes n} \to A$ and the *n*-th comultiplication $\Delta_n: C \to C^{\otimes n}$ inductively by:

 $m_0 = u, \quad m_{n+1} = m(\mathrm{id}_A \otimes m_n), \quad \Delta_0 = \varepsilon, \quad \mathrm{and} \quad \Delta_{n+1} = (\mathrm{id}_C \otimes \Delta_n)\Delta.$

For $n \geq 1$, we depict them as

$$\underline{\operatorname{PSfrag replanments}}_{A} \underbrace{\underbrace{\operatorname{PSfrag deplanments}}_{n \text{ times}} \underbrace{\operatorname{PSfrag deplanments}}_{C} \cdot \cdot \cdot$$

The (co)associativity and (co)unitality of m and Δ imply that

2.8. Categorical bialgebras. Let \mathcal{B} be a braided monoidal category. A *bialgebra* in \mathcal{B} is a quintuple $(A, m, u, \Delta, \varepsilon)$ such that (A, m, u) is an algebra in \mathcal{B} , (A, Δ, ε) is a coalgebra in \mathcal{B} , and the following compatibility relations hold:

$$\Delta m = (m \otimes m)(\mathrm{id}_A \otimes \tau_{A,A} \otimes \mathrm{id}_A)(\Delta \otimes \Delta), \quad \Delta u = u \otimes u, \quad \varepsilon m = \varepsilon \otimes \varepsilon, \quad \mathrm{and} \quad \varepsilon u = \mathrm{id}_1.$$

Graphically, these rewrite as

$$\bigcup_{i=1}^{i} = \bigcup_{i=1}^{i} \bigcup_{i=1}^{i} = \bigcup_{i=1}^{i} \bigcup_{i=1}^{i}$$

A bialgebra morphism between two bialgebras A et A' is a morphism $A \to A'$ in \mathcal{B} which is both an algebra and a coalgebra morphism.

2.9. Categorical Hopf algebras. A Hopf algebra in \mathcal{B} is a sextuple $(H, m, u, \Delta, \varepsilon, S)$, where $(H, m, u, \Delta, \varepsilon)$ is a bialgebra in \mathcal{B} and $S: H \to H$ is an isomorphism in \mathcal{B} , called the *antipode*, which satisfies

$$m(S \otimes \mathrm{id}_H)\Delta = u\epsilon = m(\mathrm{id}_H \otimes S)\Delta.$$

The antipode and its inverse are depicted by

$$S = \phi$$
 and $S^{-1} = \phi$.

Graphically, the antipode axiom is rewritten as

A useful feature of antipodes is that it is *anti-multiplicative*:

and anti-comultiplication
$$H$$
 and H and H

A *Hopf algebra morphism* between two Hopf algebras is a bialgebra morphism between them.

2.10. Categorical modules. Let (A, m, u) an algebra in a monoidal category \mathcal{B} . A left Amodule in \mathcal{B} is a pair (M, r), where $r: A \otimes M \to M$ is a morphism in \mathcal{B} , called the action of A on M, which satisfies

$$r(m \otimes \mathrm{id}_M) = r(\mathrm{id}_A \otimes r)$$
 and $r(u \otimes \mathrm{id}_M) = \mathrm{id}_M$.

Graphically, the action $r: A \otimes M \to M$ is denoted by



A morphism $f: (M, r) \to (M', r')$ between two left A-modules (M, r) and (M', r') is a morphism $f: M \to M'$ in \mathcal{B} such that $fr = r'(\mathrm{id}_A \otimes f)$. With composition inherited from \mathcal{B} , left A-modules and morphisms between them form a category ${}_A\mathbf{Mod}$.

When \mathcal{B} is braided and A is a bialgebra in \mathcal{B} , the category ${}_{A}\mathbf{Mod}$ is monoidal: the unit object of ${}_{A}\mathbf{Mod}$ is the pair $(\mathbb{1}, \varepsilon)$, the monoidal product of two left A-modules (M, r) and (M', r') is given by the pair $(M \otimes M', s)$, where

$$s = (r \otimes r')(\mathrm{id}_A \otimes \tau_{A,M} \otimes \mathrm{id}_{M'})(\Delta \otimes \operatorname{id}_{M \otimes M'}) = \bigwedge_{A = M = M'}^{M = M'},$$

and the monoidal product of morphisms is inherited from \mathcal{B} .

IVAN BARTULOVIĆ

2.11. Categorical comodules. Let (C, Δ, ε) a coalgebra in a monoidal category \mathcal{B} . A *left C-comodule* in \mathcal{B} is a pair (N, γ) , where $\gamma \colon N \to C \otimes N$ is a morphism in \mathcal{B} , called *the coaction of* C *on* N, which satisfies

$$(\Delta \otimes \mathrm{id}_N)\gamma = (\mathrm{id}_C \otimes \gamma)\gamma$$
 and $(\varepsilon \otimes \mathrm{id}_N)\gamma = \mathrm{id}_N$.

Graphically, the coaction $\gamma \colon N \to C \otimes N$ is denoted by

 $\frac{PSfrag \text{ replacements}}{\gamma =} \underbrace{\frac{PSfrag \text{ replacements}}{Sfrag \text{ replacements}}}_{PSfrag \text{ replacements}} \underbrace{\frac{PSfrag \text{ replacements}}{C}}_{N} \\ \text{so that the axioms of } \underbrace{\frac{RSfefts}{C} \cdot \frac{APleonmenta}{N}}_{C} \text{lle rewrite as } \underbrace{\frac{C}{N} \cdot \frac{N}{C}}_{N} \\ \underbrace{\frac{PSfrag \text{ replacements}}{C}}_{N} \underbrace{\frac{C}{N} \cdot \frac{N}{C}}_{N} \\ \underbrace{\frac{PSfrag \text{ replacements}}{C}}_{N} \underbrace{\frac{C}{N} \cdot \frac{N}{N}}_{N} \\ \underbrace{\frac{C}{N} \cdot \frac{N}{N}}_{N} \underbrace{\frac{C}{N} \cdot \frac{N}{N}}_{N} \\ \frac{C}{N} \cdot \frac{N}{N} \\ \underbrace{\frac{C}{N} \cdot \frac{N}{N}}_{N} \\ \frac{C}{N} \cdot \frac{N}{N} \\ \frac{C}{N} \\ \frac{C$

A morphism $f: (N, \gamma) \to (N', \gamma')$ between two left A-comodules (N, γ) and (N', γ') is a morphism $f: N \to N'$ in \mathcal{B} such that $\gamma' f = (\mathrm{id}_C \otimes f)\gamma$. With composition inherited from \mathcal{B} , left C-comodules and morphisms between them form a category $_C$ **Comod**.

When \mathcal{B} is braided and C is a bialgebra in \mathcal{B} , the category $_{C}$ **Comod** is monoidal: the unit object of $_{C}$ **Comod** is the pair $(\mathbb{1}, u)$, the monoidal product of two left C-comodules (N, γ) and (N', γ') is given by the pair $(N \otimes N', \delta)$, where

$$\delta = (m \otimes \mathrm{id}_{N \otimes N'})(\mathrm{id}_C \otimes \tau_{N,C} \otimes \mathrm{id}_{N'})(\gamma \otimes \gamma) = \bigcup_{N = N'}^{C \quad N \quad N'} ,$$

and the monoidal product of morphisms is inherited from \mathcal{B} .

2.12. **Diagonal actions.** Let H be a brace production of H on $H^{\otimes n}$ is defined inductively by $\mathbb{1}_{H}^{\mathbb{S}$ frag replacements in a brace of \mathcal{B} . The left diagonal \mathbb{R} is defined inductively by $\mathbb{1}_{H}^{\mathbb{S}}$

$$\begin{array}{c|c} \underline{\operatorname{PSfrag replacements}} & \underline{\operatorname{PSfrag replacements}} & \underline{\operatorname{H}} & \underline$$



n times

2.15. Pivotal categories. A *pivotal* category is a monoidal category \mathcal{B} such that each object X of \mathcal{B} has a *dual* object X^* and four morphisms

$$ev_X \colon X^* \otimes X \to \mathbb{1}, \quad coev_X \colon \mathbb{1} \to X \otimes X^*, \\ ev_X \colon X \otimes X^* \to \mathbb{1}, \quad \widetilde{coev_X} \colon \mathbb{1} \to X^* \otimes X,$$

satisfying some conditions. Briefly, these say that the associated left/right dual functors coincide as monoidal functors (see [12, Chapter 1] for more details). The latter implies that the dual morphism $f^* \colon Y^* \to X^*$ of a morphism $f \colon X \to Y$ in \mathcal{B} is computed by

$$f^* = (\mathrm{id}_{X^*} \otimes \widetilde{\mathrm{ev}}_Y)(\mathrm{id}_{X^*} \otimes f \otimes \mathrm{id}_{Y^*})(\widetilde{\mathrm{coev}}_X \otimes \mathrm{id}_{Y^*}) = \\ = (\mathrm{ev}_Y \otimes \mathrm{id}_{X^*})(\mathrm{id}_{Y^*} \otimes f \otimes \mathrm{id}_{X^*})(\mathrm{id}_{Y^*} \otimes \mathrm{coev}_X).$$

We extend the graphical calculus for monoidal $\overset{\text{PSfrag replacements}}{\text{categories (see Section 2.4)}}$ to pivotal categories by orienting arcs. If an arc colored by X is oriented upwards, the represented object in source/target of corresponding morphism is X^* . For example, $\mathrm{id}_X, \mathrm{id}_{X^*}$, and a morphism $f: X \otimes Y^* \otimes Z \to U \otimes V^*$ are depicted by

$$\underbrace{\frac{PSfrag \text{ replacements}}{PSfrag \text{ replacements}}}_{X} = \underbrace{\frac{PSfrag \text{ replacements}}{X}}_{X} = \underbrace{\frac{X}{X}}_{X}, \text{ and } f = \underbrace{\frac{\Psi U \Psi V}{V}}_{X \Psi Z}.$$
The morphisms $ev_{X} \underbrace{ev_{X}, co \underbrace{PSfrag \text{ replacements}}_{X} \underbrace{PSfrag \text{ replacements}}_{X}}_{X} = \underbrace{\frac{PSfrag \text{ replacements}}_{X}}_{X} = \underbrace{\frac{PSfrag \text{ replacements}}_{X}}_{X}, \underbrace{\frac{PSfrag \text{ replacements}}_{X}}_{X} = \underbrace{\frac{V \Psi V}{V}}_{X}.$

2.16. Left and right twists. Let \mathcal{B} be a braided pivotal category. The *left twist* of an object X of \mathcal{B} is defined by

$$\underline{\operatorname{PSfrag repla}}_{X} \stackrel{l}{\underset{X}{\operatorname{perfrag repla}}} \bigvee_{X} = (\operatorname{id}_{X} \otimes \widetilde{\operatorname{ev}}_{X})(\tau_{X,X} \otimes \operatorname{id}_{X^{*}})(\operatorname{id}_{X} \otimes \operatorname{coev}_{X}) \colon X \to X,$$

while the *right twist* of X is defined by

ī

$$\underline{\operatorname{PSfrag replater reprint}}_{X} \bigcirc \bigvee_{X} = (\operatorname{ev}_{X} \otimes \operatorname{id}_{X})(\operatorname{id}_{X^{*}} \otimes \tau_{X,X})(\widetilde{\operatorname{coev}}_{X} \otimes \operatorname{id}_{X}) \colon X \to X.$$

The left and the right twist are natural isomorphisms with inverses

$$\underbrace{\operatorname{PSfrag}\left(\theta_{\text{pta}}^{l}\right)^{-1}_{\text{cements}}}_{X} \qquad \underbrace{\operatorname{PSfrag}\left(\theta_{X}^{l}\right)^{-1}}_{X} \xrightarrow{\mathbb{P}}_{X} \xrightarrow{\mathbb{P}}_{X$$

The left twist $\theta^l = \{\theta_X^l \colon X \to X\}_{X \in Ob(\mathcal{B})}$ and the right twist $\theta^r = \{\theta_X^r \colon X \to X\}_{X \in Ob(\mathcal{B})}$ are twists for \mathcal{B} in the sense of Section 2.3.

A ribbon category is a braided pivotal category \mathcal{B} whose left and right twist coincide. Then $\theta = \theta_l = \theta_r$ is called the *twist* of \mathcal{B} .

2.17. Coends. Let \mathcal{C} and \mathcal{D} be any categories and $F: \mathcal{C}^{\text{op}} \times \mathcal{C} \to \mathcal{D}$ a functor. A *dinatural* transformation between F and an object D in \mathcal{D} is a function d that assigns to any object X in \mathcal{C} a morphism $d_X: F(X, X) \to D$ such that for all morphisms $f: X \to Y$ in \mathcal{C}

the following diagram commutes:



A coend of a functor $F: \mathcal{C}^{\text{op}} \times \mathcal{C} \to \mathcal{D}$ is a pair (C, i) where C is an object of \mathcal{D} and i is a dinatural transformation from F to C, which is universal among all dinatural transformations. More precisely, for any dinatural transformation d from F to D, there exists a unique morphism $\varphi: C \to D$ in \mathcal{D} such that $d_X = \varphi i_X$ for all $X \in \text{Ob}(\mathcal{C})$. A coend (C, i) of a functor F, if it exists, is unique up to a unique isomorphism commuting with the dinatural transformation.

2.18. Coend of a pivotal category. Let \mathcal{B} be a pivotal category. The *coend* of \mathcal{B} , if it exists, is the coend (H, i) of the functor $F: \mathcal{B}^{\text{op}} \times \mathcal{B} \to \mathcal{B}$ defined by

$$F(X,Y) = X^* \otimes Y$$
 and $F(f,g) = f^* \otimes g$

YH

We depict the universal dinatural transformation $i \stackrel{\text{PSfrag replacements}}{\text{PSfrag replacements}} \to H_{X \in Ob(\mathcal{B})}$ as

$$i_X = \begin{array}{c} H \\ X \\ Y \\ Y \\ X \\ H \end{array}$$

Note that H is a coalgebra $\stackrel{X}{H} \mathcal{B}$ with comultiplication $\Delta_X^{\Delta} H \to H \otimes_X^{\mathcal{H}} H$ and counit $\varepsilon \colon H \to \mathbb{1}$, which are unique morphisms such that, for all $X \in Ob(\mathcal{B})$, X

The coalgebra (H, Δ, ε) coacts on the objects in \mathcal{B} via the *universal coaction* defined for any $X \in Ob(\mathcal{B})$ by

$$\delta_X = (\mathrm{id}_X \otimes i_X) \circ (\mathrm{coev}_X \otimes \mathrm{id}_X).$$

We will denote it graphically as

$$\begin{array}{c|c} PS \text{frag replacements} \\ X \end{array} \xrightarrow{PS \text{frag replacements}} \\ X \end{array} X$$

Note that δ_H is the right coadjoint coaction of H on H (see Section 2.14). If \mathcal{B} is braided, then H is a Hopf algebra in \mathcal{B} . Its unit is $u = \delta_1 : \mathbb{1} \xrightarrow{X} H$ and \inf_{H}^m multiplication $m : H \otimes H \to H$ and antipode \mathbb{P} fragge place preserve characterized as follows: for fall $X, Y \in Ob(\mathcal{B})$,



We refer to [12, Chapter 6] for details.

3. SIMPLICIAL, PARACYCLIC, AND CYCLIC OBJECTS

In this section we recall the notions of (co)simplicial, para(co)cyclic, and (co)cyclic objects in a category.

IVAN BARTULOVIĆ

3.1. The simplicial category. The simplicial category Δ is defined as follows. The objects of Δ are the nonnegative integers $n \in \mathbb{N}$. A morphism $n \to m$ in Δ is an increasing map between sets $[n] = \{0, \ldots, n\}$ and $[m] = \{0, \ldots, m\}$. For $n \in \mathbb{N}^*$ and $0 \leq i \leq n$, the *i*-th coface $\delta_i^n : n - 1 \to n$ is the unique increasing injection from [n - 1] into [n] which misses $i \in [n]$. For $n \in \mathbb{N}$ and $0 \leq j \leq n$, the *j*-th codegeneracy $\sigma_j^n : n + 1 \to n$ is the unique increasing surjection from [n + 1] onto [n] which sends both j and j + 1 to j.

It is well known (see [10, Lemma 5.1]) that morphisms in Δ are generated by cofaces $\{\delta_i^n\}_{0 \le i \le n, n \in \mathbb{N}^*}$ and codegeneracies $\{\sigma_j^n\}_{0 \le j \le n, n \in \mathbb{N}}$ subject to the simplicial relations (SR):

$$\delta_{j}\delta_{i} = \delta_{i}\delta_{j-1} \quad \text{for } i < j,$$

$$\sigma_{j}\sigma_{i} = \sigma_{i}\sigma_{j+1} \quad \text{for } i \leq j,$$

(SR)
$$\sigma_{j}\delta_{i} = \begin{cases} \delta_{i}\sigma_{j-1} & \text{for } i < j, \\ \text{id}_{n} & \text{for } i = j, i = j+1, \\ \delta_{i-1}\sigma_{j} & \text{for } i > j+1. \end{cases}$$

3.2. The paracyclic category. The paracyclic category ΔC_{∞} is defined as follows. The objects of ΔC_{∞} are the nonnegative integers $n \in \mathbb{N}$. The morphisms are generated by morphisms $\{\delta_i^n\}_{n \in \mathbb{N}^*, 0 \leq i \leq n}$, called *cofaces*, morphisms $\{\sigma_j^n\}_{n \in \mathbb{N}, 0 \leq j \leq n}$, called *codegeneracies*, and isomorphisms $\{\tau_n : n \to n\}_{n \in \mathbb{N}}$, called *paracocyclic operators*, satisfying the simplicial relations (SR) and the following *paracyclic compatibility relations* (PCR):

(PCR)
$$\begin{aligned} \tau_n \delta_i &= \delta_{i-1} \tau_{n-1} \quad \text{for } 1 \leq i \leq n, \\ \tau_n \delta_0 &= \delta_n, \\ \tau_n \sigma_i &= \sigma_{i-1} \tau_{n+1} \quad \text{for } 1 \leq i \leq n, \text{ and} \\ \tau_n \sigma_0 &= \sigma_n \tau_{n+1}^2. \end{aligned}$$

Note that Δ is a subcategory of ΔC_{∞} .

3.3. The cyclic category. The cyclic category ΔC is defined as follows. The objects of ΔC are the nonnegative integers $n \in \mathbb{N}$. The morphisms are generated by morphisms $\{\delta_i^n\}_{n\in\mathbb{N}^*,0\leq i\leq n}$, called cofaces, morphisms $\{\sigma_j^n\}_{n\in\mathbb{N},0\leq j\leq n}$, called codegeneracies, and isomorphisms $\{\tau_n: n \to n\}_{n\in\mathbb{N}}$, called cocyclic operators, which satisfy the relations (SR), (PCR), and the cyclicity condition (CC):

(CC)
$$\tau_n^{n+1} = \mathrm{id}_n.$$

Note that ΔC is a quotient of ΔC_{∞} .

3.4. (Co)simplicial, para(co)cyclic, and (co)cyclic objects in a category. Let \mathcal{C} be any category. A simplicial object in \mathcal{C} is a functor $X : \Delta^{\text{op}} \to \mathcal{C}$, a paracyclic object in \mathcal{C} is a functor $\Delta C_{\infty}^{\text{op}} \to \mathcal{C}$, and a cyclic object in \mathcal{C} is a functor $\Delta C^{\text{op}} \to \mathcal{C}$. Dually, a cosimplicial object in \mathcal{C} is a functor $\Delta \to \mathcal{C}$, a paracocyclic object in \mathcal{C} is a functor $\Delta C_{\infty} \to \mathcal{C}$, and a cocyclic object in \mathcal{C} is a functor $\Delta \to \mathcal{C}$, a paracocyclic object in \mathcal{C} is a functor $\Delta C_{\infty} \to \mathcal{C}$, and a cocyclic object in \mathcal{C} is a functor $\Delta C \to \mathcal{C}$. A (co)simplicial/para(co)cyclic/(co)cyclic object in the category of sets (respectively, of k-modules) are called (co)simplicial/para(co)cyclic/(co)cyclic sets (respectively, k-modules).

A morphism between two (co)simplicial/para(co)cyclic/(co)cyclic objects is a natural transformation between them. One often denotes the image of a morphism f under a (co)simplicial/para(co)cyclic/(co)cyclic by the same letter f.

Since the categories $\Delta, \Delta C_{\infty}, \Delta C$ are defined by generators and relations, a (co)simplicial/para(co)cyclic/(co)cyclic object in a category is entirely determined by the images of the generators satisfying the corresponding relations. For example, a paracocyclic object Y in \mathcal{C} may be seen as a family $Y_{\bullet} = \{Y_n\}_{n \in \mathbb{N}}$ of objects of \mathcal{C} equipped with morphisms $\{\delta_i^n \colon Y_{n-1} \to Y_n\}_{n \in \mathbb{N}^*, 0 \leq i \leq n}$, called *cofaces*, morphisms $\{\sigma_j^n \colon Y_{n+1} \to Y_n\}_{n \in \mathbb{N}, 0 \leq j \leq n}$, called *codegeneracies*, and isomorphisms $\{\tau_n \colon Y_n \to Y_n\}_{n \in \mathbb{N}}$, called *paracocyclic operators*, subject to the relations (SR) and (PCR). Similarly, a morphism $\beta \colon Y \to Y'$ between two paracocyclic objects Y_{\bullet} and Y'_{\bullet} in \mathcal{C} is explicited as a family $\{\beta_n \colon Y_n \to Y'_n\}_{n \in \mathbb{N}}$ of morphisms in \mathcal{C} satisfying

$$\beta_n \delta_i^n = \delta_i^n \beta_{n-1} \quad \text{for any } n \in \mathbb{N}^* \text{ and } 0 \le i \le n,$$

$$\beta_n \sigma_j^n = \sigma_j^n \beta_{n+1} \quad \text{for any } n \in \mathbb{N} \text{ and } 0 \le j \le n,$$

$$\beta_n \tau_n = \tau_n \beta_n \quad \text{for any } n \in \mathbb{N}.$$

Clearly, the composition of a (co)simplicial/para(co)cyclic/(co)cyclic object X in \mathcal{C} with a functor $F: \mathcal{C} \to \mathcal{D}$ is a (co)simplicial/para(co)cyclic/(co)cyclic object FX in \mathcal{D} . In particular, useful examples are provided by the covariant and the contravariant Hom-functors $\operatorname{Hom}_{\mathcal{C}}(I, -)$ and $\operatorname{Hom}_{\mathcal{C}}(-, I)$, where I is an object of \mathcal{C} . In this case, we denote:

 $\operatorname{Hom}_{\mathcal{C}}(I, X) = \operatorname{Hom}_{\mathcal{C}}(I, -) \circ X$ and $\operatorname{Hom}_{\mathcal{C}}(X, I) = \operatorname{Hom}_{\mathcal{C}}(-, I) \circ X$.

Lemma 1. Let \mathcal{B} be a k-linear braided category with a twist θ . If $Y_{\bullet} = \{Y_n\}_{n \in \mathbb{N}}$ is a paracocyclic object in \mathcal{B} such that its paracocyclic operator satisfies $\tau_n^{n+1} = \theta_{Y_n}$ for each $n \in \mathbb{N}$, then

- (a) $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_{\bullet}) = \{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n)\}_{n \in \mathbb{N}} \text{ is a cocyclic } \mathbb{k}\text{-module},$
- (b) $\operatorname{Hom}_{\mathcal{B}}(Y_{\bullet}, \mathbb{1}) = {\operatorname{Hom}_{\mathcal{B}}(Y_n, \mathbb{1})}_{n \in \mathbb{N}}$ is a cyclic k-module.

Proof. Let us prove (a). By composition with $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, -) \colon \mathcal{B} \to \operatorname{Mod}_{\Bbbk}$, we obtain that $\{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n)\}_{n \in \mathbb{N}}$ is a paracocyclic \Bbbk -module. Let us now verify that (CC) holds. Let $n \in \mathbb{N}$. The morphism $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, -)(\theta_{Y_n}) \colon \operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n) \to \operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n)$ is the \Bbbk -linear morphism given by $f \mapsto \theta_{Y_n} f$. The naturality of θ and the fact that $\theta_{\mathbb{1}} = \operatorname{id}_{\mathbb{1}}$ imply that $\theta_{Y_n} f = f \theta_{\mathbb{1}} = f$ for all $f \in \operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n)$. Then, using the functoriality of Y_{\bullet} and the hypothesis that $Y_{\bullet}(\tau_n^{n+1}) = \theta_{Y_n}$, we have

$$\left(\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_{\bullet})(\tau_n)\right)^{n+1} = \operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, -)\left(Y_{\bullet}(\tau_n^{n+1})\right) = \operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, -)(\theta_{Y_n}) = \operatorname{id}_{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, Y_n)}.$$

Part (b) is proved similarly.

3.5. Cyclic (co)homology. To any cyclic k-module $X: \Delta C^{\text{op}} \to \text{Mod}_{\Bbbk}$, one can associate a bicomplex CC(X) (see [13]). The *n*-th cyclic homology $HC_n(X)$ of X is defined as the *n*-th homology of the total chain complex associated to the chain bicomplex CC(X). A morphism between cyclic k-modules induces a levelwise morphism in cyclic homology.

Similarly, to any cocyclic k-module $Y: \Delta C \to \mathbf{Mod}_k$, one can associate a cochain bicomplex $\mathbf{CC}(Y)$, obtained by a construction dual to the one of a chain bicomplex. The *n*-th cyclic cohomology $HC^n(Y)$ of Y is defined as the *n*-th cohomology of the total cochain complex associated to the cochain bicomplex $\mathbf{CC}(Y)$. A morphism between cocyclic k-modules induces a levelwise morphism in cyclic cohomology.

4. MODULAR PAIRS AND BRAIDED CONNES-MOSCOVICI CONSTRUCTION

In this section, \mathcal{B} is a braided monoidal category and H is a Hopf algebra in \mathcal{B} . We provide a braided generalization of the notion of a modular pair in involution for H and then compute the powers of the paracocyclic operator associated to such a pair (see Theorem 2 and its corollaries).

IVAN BARTULOVIĆ

4.1. Modular pairs. A modular pair for H is a pair (δ, σ) where $\delta: H \to \mathbb{1}$ is an algebra morphism and $\sigma: \mathbb{1} \to H$ is a coalgebra morphism such that $\delta\sigma = \mathrm{id}_{\mathbb{1}}$. For instance, (ε, u) is a modular pair for H, where $\varepsilon: H \to \mathbb{1}$ and $u: \mathbb{1} \to H$ are the counit and unit of H, respectively.

Given a twist θ for \mathcal{B} , a θ -twisted modular pair in involution for H is a modular pair (δ, σ) for H such that



Here we use the graphical conventions from Section 2.4. If H is *involutive* Hopf algebra in \mathcal{B} in the sense that $S^2 = \theta_H$, then (ε, u) is a θ -twisted modular pair in involution for H.

Note that if \mathcal{B} is symmetric with a trivial twist $\mathrm{id}_{\mathcal{B}}$ (see Section 2.3), then an $\mathrm{id}_{\mathcal{B}}$ -twisted modular pair in involution corresponds to a braided modular pair in involution in the sense of [8].

4.2. Powers of the paracocyclic operator $\tau_n(\delta,\sigma)$: $H_{H^{\otimes n-1}}^{\otimes n}$ be a modular pair for H. For any $n \geq 0$, define the paracocyclic operator $\tau_n(\delta,\sigma)$: $H_{H^{\otimes n-1}}^{\otimes n} H^{\otimes n}$ by

$$\tau_0(\delta,\sigma) = \mathrm{id}_{\mathbbm{1}}, \quad \tau_1(\delta,\sigma_{\!H}) \stackrel{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}}{\underset{H}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{\overset{\mathfrak{S}_{\mathrm{sm-1}}}}{{}}}}}}}}}}}}}}}}}}}}}}}}}}$$

Here we use the diagonal actions defined in Section 2.12. Note that the operators $\tau_n(\delta, \sigma)$ are the paracocyclic operators of a paracocyclic object in \mathcal{B} associated with H and (δ, σ) (see Section 4.3). In the following theorem we compute the powers (up to n+1) of $\tau_n(\delta, \sigma)$.

Theorem 2. For $n \ge 2$ and $2 \le k \le n$, we have:

(4)
$$\frac{PSfrag \ replacements}{\left(\tau_{n}(\delta,\sigma)\right)^{k}} = \underbrace{\tau_{1}(\delta,\sigma)}_{\substack{H^{\otimes k-2} \ k-1 \ k}} \underbrace{\tau_{n}(\varepsilon,u)^{k-1}}_{\substack{\delta \\ K^{\otimes k-2} \ k-1}} \underbrace{\tau_{n}(\varepsilon,u)^{k-1}}_{\substack{\delta \\ K^{\otimes k-2} \ k-1$$

In addition,

(5)
$$\underbrace{PSfrag \ replacements}_{H \ H} (\tau_n(\delta, \sigma))^{n+1} = \begin{cases} 0 & \text{if } n = 1, \\ H & H \\ H^{\otimes n-1} & \text{or} \\ \hline (\tau_1(\delta, \sigma))^2 & (\tau_{n-1}(\varepsilon, u))^n \\ \hline (\tau_{n-1}(\varepsilon, u))^n & \text{if } n \ge 2. \end{cases}$$

In the statements of the theorem, we use the diagonal actions together with the (co)adjoint (co)actions defined in Sections 2.13 and 2.14. Also, an integer k below an arc denotes the k-th tensorand of $H^{\otimes n}$. We prove Theorem 2 in Section 6 by induction and by using properties of modular pairs and twisted antipodes.

In the next corollary, we compute the (n + 1)-th power of the paracocyclic operator $\tau_n(\delta, \sigma)$ in terms of the (n + 1)-th power of the paracocyclic operator $\tau_n(\varepsilon, u)$, where ε and u are the counit and unit of H.

Corollary 3. For any modular pair (δ, σ) and any $n \in \mathbb{N}$,



Proof. We show the result by induction. For n = 0, this follows since the adjoint action on $\mathbb{1}$ is given by counit, since the coadjoint coaction on $\mathbb{1}$ is given by unit and the fact that $\tau_0(\varepsilon, u) = \tau_0(\delta, \sigma) = \mathrm{id}_{\mathbb{1}}$. Let us check the case n = 1. Using (5) and the fact that $\tau_1(\varepsilon, u) = S$, we obtain





Here (i) follows by applying (5) for the modular pair (δ, σ) , (ii) by applying the result for n = 1, and (iii) by applying (5) for the modular pair (ε, u) .

The next corollary states that the paracocyclic operator associated with a twisted modular pair in involution satisfies the "twisted cocyclicity condition".

Corollary 4. If \mathcal{B} has a twist θ and (δ, σ) is a θ -twisted modular pair in involution for H, then $(\tau_n(\delta, \sigma))^{n+1} = \theta_{H^{\otimes n}}$ for all $n \in \mathbb{N}$.

Proof. The equality $(\tau_n(\delta, \sigma))^{n+1} = \theta_{H^{\otimes n}}$ is shown by induction. For n = 0, this follows by definition and the fact that $\theta_1 = \frac{1}{PSHdg} =$



Here (i) follows by Formula (5) of Theorem 2, (ii) follows ${}^{\tau}$ by the fact that (δ, σ) is a θ -twisted modular pair in involution for H, (iii) follows by the naturality of the twist and the definitions of left coadjoint coaction and right adjoint action, (iv) follows by (co)associativity and the fact that Φ^{PStrag} replacements algebra morphism and $H \delta^n$ is a coalgebra morphism, (v) follows by the antipode axion $H^{\otimes n}_{\text{and}}$ (co)unitality. $H^{\otimes n}_{H}$

Suppose that the statement is true for $n \ge 1$ and let u_{π} by it_{1}^{H} for n+1. We have



Here (i) follows from Formula (5) of Theorem 2, (ii) follows from Corollary 3, (iii) follows from the statement for n = 1 and the induction hypothesis, (iv) follows by the naturality of the braiding and from the axiom of the twist.

Remark 5. If \mathcal{B} is a symmetric monoidal category endowed with the trivial twist $\mathrm{id}_{\mathcal{B}}$ and (δ, σ) is a $\mathrm{id}_{\mathcal{B}}$ -twisted modular pair for H, then Corollary 4 gives that $(\tau_n(\delta, \sigma))^{n+1} =$ $\mathrm{id}_{H^{\otimes n}}$ for all $n \in \mathbb{N}$. This was first proved by Khalkhali and Pourkia in [8]. 4.3. Paracocyclic objects associated with modular pairs. Let (δ, σ) be a modular pair for H. Let us recall the paracocyclic object $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ in \mathcal{B} from [8] associated to this data. For any $n \geq 0$, define

$$\mathbf{CM}_n(H,\delta,\sigma) = H^{\otimes n}.$$

For any $n \ge 1$, define the cofaces $\{\delta_i^n(\sigma) \colon H^{\otimes n-1} \to H^{\otimes n}\}_{0 \le i \le n}$ by setting $\delta_0^1 = u$, $\delta_1^1 = \sigma$, and for any $n \ge 2$,

For any $n \ge 0$, define the codegeneracies $\{\sigma_j^n \colon H^{\otimes n+1} \to H^{\otimes n}\}_{0 \le j \le n}$ by

$$\frac{\operatorname{PSfrag replacements}}{\sigma_j^n} = \left| \cdots \right| \quad \begin{pmatrix} \mathsf{O} \\ \mathsf{O} \\ \mathsf{I} \\ j+1 \\ n+1 \end{pmatrix} .$$

For any $n \ge 0$, the paracocyclic operators $\tau_n(\delta, \sigma) \colon H^{\otimes n} \to H^{\otimes n}$ of $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ are those defined in Section 4.2.

Theorem 2 is useful to prove that $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ is a paracocyclic object in \mathcal{B} . We prove this in Appendix. In particular, we prove that for all $n \in \mathbb{N}$,

$$\tau_n(\delta,\sigma)\sigma_0^n = \sigma_n^n(\tau_{n+1}(\delta,\sigma))^2.$$

The next corollary derives (co)cyclic k-modules from $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$. It follows directly from Lemma 1 and Corollary 4.

Corollary 6. If \mathcal{B} has a twist θ and (δ, σ) a θ -twisted modular pair in involution for H, then

- (a) $\{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \operatorname{\mathbf{CM}}_n(H, \delta, \sigma))\}_{n \in \mathbb{N}}$ is a cocyclic k-module,
- (b) $\{\operatorname{Hom}_{\mathcal{B}}(\mathbf{CM}_n(H, \delta, \sigma), \mathbb{1})\}_{n \in \mathbb{N}}$ is a cyclic k-module.

5. CATEGORICAL CONNES-MOSCOVICI TRACE

In this section, \mathcal{B} is a braided category with a twist θ and H is a Hopf algebra in \mathcal{B} . We introduce traces (à la Connes-Moscovici) between paracocyclic objects associated with H (as in Section 4.3) and paracocyclic objects associated with an H-module coalgebra. We provide an explicit example of such traces using coends.

IVAN BARTULOVIĆ

5.1. Paracocyclic objects associated with coalgebras. Let C be a coalgebra in \mathcal{B} . We associate with C a paracocyclic object $C_{\bullet}(C)$ in \mathcal{B} . It is inspired by the construction of Akrami and Majid from [1]. When $\mathcal{B} = \mathbf{Mod}_{\Bbbk}$ is the category of \Bbbk -modules, one recovers the cocyclic k-module implicitly defined in the work of Farinati and Solotar [7]. When \mathcal{B} is a symmetric monoidal category endowed with the trivial twist, then the underlying cosimplicial object of $\mathbf{C}_{\bullet}(C)$ is equal to the one considered in [2, Definition 2.2].

For $n \ge 0$, set $\mathbf{C}_n(C) = C^{\otimes n+1}$. For $n \ge 1$, define the cofaces $\{\delta_i^n : C^{\otimes n} \to C^{\otimes n+1}\}_{0 \le i \le n}$ by

For $n \ge 0$, define the codegeneracies $\{\sigma_j^n \colon C^{\otimes n+2} \to C^{\otimes n+1}\}_{0 \le j \le n}$ by

$$\frac{\operatorname{PSfrag replacements}}{\sigma_j^n} = \left| \cdots \right| \left| \begin{array}{c} \mathsf{O} \\ \mathsf{O} \end{array} \right| \left| \cdots \right| .$$

$$0 \qquad j+1 \qquad n+1$$

For $n \ge 0$, define the paraces by extreme paraces $\tau_n : C^{\otimes n+1} \to C^{\otimes n+1}$ by

$$\tau_0 = \bigcirc_C \quad \text{and} \quad \tau_n = \bigcirc_{\substack{C \\ n-1 \\ 0 \\ 1 \\ n}} \quad \text{if} \quad n \ge 1.$$

It follows directly from the definition of a twist (see Section 2.3) that the paracocyclic operator for $\mathbf{C}_{\bullet}(C)$ satisfies the relation $\tau_n^{n+1} = \theta_{C^{\otimes n+1}}$ for all $n \in \mathbb{N}$.

5.2. Traces. Let C be an H-module coalgebra in \mathcal{B} , that is, a coalgebra in the category of right H-modules in \mathcal{B} . In other words, C is a coalgebra in \mathcal{B} endowed with a right action $r: C \otimes H \to C$ of H on C such that the comultiplication Δ_C and the counit ε_C of C are both H-linear, that is, morphisms of right H-modules. By depicting the right action by



In this pictures, the red strands are colored by C and the black ones by H.

Let $\delta: H \to \mathbb{1}$ be an algebra morphism and let $\sigma: \mathbb{1} \hookrightarrow H$ be a coalgebra morphism. A δ -invariant σ -trace for C is a morphist $\alpha \colon \mathbb{1} \to C$ in \mathcal{B} satisfying



Given such a morphism, define for any $n \in \mathbb{N}$ the morphism $\alpha_n \colon H^{\otimes n} \to C^{\otimes n+1}$ in \mathcal{B} by setting



Consider the paracocyclic object $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ in \mathcal{B} (see Section 4.3) and the paracocyclic object $\mathbf{C}_{\bullet}(C)$ in \mathcal{B} associated to the coalgebra C in \mathcal{B} (see Section 5.1).

Theorem 7. Let (δ, σ) be a modular pair for H and α be a δ -invariant σ -trace for C. Then the family $\{\alpha_n \colon H^{\otimes n} \to C^{\otimes n+1}\}_{n \in \mathbb{N}}$ is a morphism of paracocyclic objects from $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ to $\mathbf{C}_{\bullet}(C)$.

We prove Theorem 7 in Section 7. The next corollary relates the cyclic (co)homologies associated with $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ and $\mathbf{C}_{\bullet}(C)$.

Corollary 8. Assume that \mathcal{B} is k-linear and has a twist θ . Let (δ, σ) be a θ -twisted modular pair in involution for H and α be a δ -invariant σ -invariant trace for C. Then

(a) The family $\{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \alpha_n)\}_{n \in \mathbb{N}}$ induces a morphism in cyclic cohomology

$$\alpha^* \colon HC^* \left(\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{CM}_{\bullet}(H, \delta, \sigma)) \right) \to HC^* \left(\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{C}_{\bullet}(C)) \right)$$

(b) The family $\{\operatorname{Hom}_{\mathcal{B}}(\alpha_n, \mathbb{1})\}_{n \in \mathbb{N}}$ induces a morphism in cyclic homology

$$\alpha_* \colon HC_* \left(\operatorname{Hom}_{\mathcal{B}}(\mathbf{C}_{\bullet}(C), \mathbb{1}) \right) \to HC_* \left(\operatorname{Hom}_{\mathcal{B}}(\mathbf{CM}_{\bullet}(H, \delta, \sigma), \mathbb{1}) \right)$$

Proof. Since $(\mathbf{C}_{\bullet}(C)(\tau_n))^{n+1} = \theta_{C^{\otimes n+1}}$, Lemma 1 implies that $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{C}_{\bullet}(C))$ is a cocyclic k-module and that $\operatorname{Hom}_{\mathcal{B}}(\mathbf{C}_{\bullet}(C), \mathbb{1})$ is a cyclic k-module. Next, by an application of Theorem 7, the family $\{\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \alpha_n)\}_{n \in \mathbb{N}}$ is a natural transformation between cocyclic kmodules $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{CM}_{\bullet}(H, \delta, \sigma))$ and $\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{C}_{\bullet}(C))$. This means that for any $n \in \mathbb{N}$, there is a morphism

$$\alpha^{n} \colon HC^{n} \left(\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{CM}_{\bullet}(H, \delta, \sigma)) \right) \to HC^{n} \left(\operatorname{Hom}_{\mathcal{B}}(\mathbb{1}, \mathbf{C}_{\bullet}(C)) \right)$$
$$[f] \mapsto [\alpha_{n} f],$$

where [f] is a representative class of an *n*-th cyclic cocycle. This finishes the proof of the part (*a*). The proof of (*b*) is similar.

5.3. Traces from coends. Let \mathcal{B} be a ribbon category with a coend (H, i), see Section 2.18. By [12, Chapter 6], the object H is a Hopf algebra in \mathcal{B} which is involutive (that is $S^2 = \theta_H$, where θ is the twist of \mathcal{B} , see Section 2.16). By Section 4.1, since H is involutive, the pair (ε, u) is a θ -twisted modular pair in involution for H. Since \mathbf{Mod}_H is braided isomorphic to the center of \mathcal{B} (see [12, Section 6.5.3] for details), we obtain that \mathbf{Mod}_H is ribbon. The braiding of \mathbf{Mod}_H is given by



Here, the coaction denoted with a black dot is the universal coaction of H (see Section 2.18). The dual of $(M, r) \in Ob(\mathbb{N}$ for the product of (M^*, r^{\dagger}) , where

$$\xrightarrow{\text{PSfrag replacements}} (\begin{array}{c} & & M \\ \hline & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & &$$

together with the (co)evaluation morphisms inherited from \mathcal{B} :

 $\operatorname{ev}_{(M,r)} = \operatorname{ev}_M, \quad \widetilde{\operatorname{ev}}_{(M,r)} = \widetilde{\operatorname{ev}}_M, \quad \operatorname{coev}_{(M,r)} = \operatorname{coev}_M, \quad \widetilde{\operatorname{coev}}_{(M,r)} = \widetilde{\operatorname{coev}}_M.$

Note that the last equality in the definition of r^{\dagger} follows from the involutivity of H.

The category \mathbf{Mod}_H has a coend ((C, a), j), where $C = H^* \otimes H$, the action $a: C \otimes H \to C$ of H on C is computed by



and the universal dinatural transformation $j = \{j_{(M,r)} \colon (M,r)^* \otimes (M,r) \to (C,a)\}_{(M,r) \in \mathbf{Mod}_H}$ is given by



By Section 2.18, the coend C is a Hopf algebra in \mathbf{Mod}_H . In particular, it is a coalgebra in \mathbf{Mod}_H . The comultiplication $\Delta_C \colon C \to C \otimes C$ and the counit $\varepsilon_C \colon C \to \mathbb{1}$ of C are computed by



The following lemma gives a way to produce an ε -invariant *u*-trace $\mathbb{1} \to C$, where ε and *u* denote the counit and unit of *H*.

Lemma 9. If a morphism $\kappa : 1 \xrightarrow{PSfrag replacements}{H in \mathcal{B} satisfies}$

$$\frac{PSfrag \ replacements}{\sigma \ \mathcal{K}} \begin{array}{c} H \\ \sigma \\ \sigma \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \\ \mathcal{K} \end{array} + \begin{array}{c} H \\ \mathcal{K} \\ \mathcal{K}$$

then the morphism

$$\underline{PSfrag \ replacements} \bigoplus^{H} \bigoplus^{H} \underset{\mathcal{K}}{\stackrel{\bullet}{\longrightarrow}} \mathbb{1} \to C$$

is an ε -invariant u-trace.

Proof. Denote $f = \varepsilon^* \otimes \operatorname{id}_H \colon H \to C$. Let us first check that f is a morphism between right H-modules (H, ε) and (C, a). Indeed,



Here (i) follows by definition of f and a, (ii) follows from the isotopy of the graphical calculus for pivotal categories and the multiplicativity of the counit, (iii) by the naturality of the braiding, the fact that $\varepsilon S = \varepsilon$, and the counitality, (iv) since universal coaction of H on itself is the right coadjoint coaction, (v) by the naturality of the braiding, the (co)unitality and the antipode axiom, (vi) follows by definition of f.

Next, $f: H \to C$ is a coalgebra morphism. Indeed, we have:



Here (i) and (v) follow by definition of f, (ii) from the isotopy of the graphical calculus for pivotal categories and the multiplicativity of the counit, (iii) by the naturality of the braiding and the fact that universal coaction of H on the state of the right coadjoint coaction, (iv) follows by the naturality of the braiding, the (co) unitality, and the antipode axiom. Also, $\frac{PSfrag replacements}{f}$



Here (i) follow from definitions of f and ε_C and (ii) from the fact that $\varepsilon u = \mathrm{id}_1$.



Here (i) and (vi) follow from definition, (ii) and (v) follow by the fact that $f: H \to C$ is a coalgebra morphism, (iii) follows by the naturality of twist and the braiding, and (iv) follows by hypothesis on κ .

Any coalgebra morphism $\mathbb{1} \to H$ satisfies the condition of Lemma 9. Another family of examples satisfying the condition of Lemma 9 is given as follows: for any $X \in Ob(\mathcal{B})$, set <u>PSfrag replacements</u>



Here, (i) and (vi) follow by definition of κ^X , (ii) and (v) follow by definition of comultiplication of H, (iii) by the naturality of twists, and (iv) by the naturality of the braiding and isotopy invariance of graphical calculus.

Thus κ^X satisfies the condition of Lemma 9 and so

(6)
$$\underline{\operatorname{PSfrag replacements}}^{X} = \bigcup_{X}^{H} : \mathbb{1} \to C$$

is an ε -invariant *u*-trace for *C*.

Remark 10. Note that if \mathcal{B} is k-linear, then any linear combination of δ -invariant σ -traces is a δ -invariant σ -trace. In particular, an interesting example of an ε -invariant u-trace comes from topological field theory: if \mathcal{B} is a ribbon fusion k-linear category and I is a representative set of simple objects of \mathcal{B} , then $\alpha = \sum_{k \in I} \dim(k) \alpha^k$ is an ε -invariant u-trace. Here α^k is defined in (6) and $\dim(k) = \widetilde{ev}_k \operatorname{coev}_k = \operatorname{ev}_k \widetilde{\operatorname{coev}}_k$ is the dimension of k.

6. Proof of Theorem 2

Our strategy to compute the (n + 1)-th power of the paracocyclic operator $\tau_n(\delta, \sigma)$ is similar to the proof of cocyclicity condition from Connes and Moscovici in [6], where Hopf algebras over \mathbb{C} are considered. We indeed proceed by induction. The difficulty here is that the paracocyclic operators involve the braiding. In our approach, based on graphical calculus, we manage to keep track the powers of paracocyclic operators. In Section 6.1 we list algebraic properties used in our proof of the equalities from Theorem 2. In Section 6.2 we show Formula (4). In Section 6.3 we show Formula (5).

Recall that H denotes a Hopf algebra in the braided monoidal category $\mathcal{B}, \delta: H \to \mathbb{1}$ is an algebra morphism and $\sigma: \mathbb{1} \to H$ is a coalgebra morphism such that $\delta \sigma = \mathrm{id}_{\mathbb{1}}$. Given such a pair, we define the *twisted antipode* $\tilde{S}: H \to H$ by



For brevity, we denote the twisted antipode $\tilde{S} \xrightarrow{\text{PSfrag replacements}}_{\text{graphically by}} \bigotimes_{H}^{\circ}$. With this notation, we will rewrite



Similarly, equation (4), which is to be proven, rewrites as



6.1. **Preliminary facts.** In this section we state several lemmas, which are used in the proof of Theorem 2. We mention that equalities (a) and (b) from Lemma 11 and the equality from Remark 17 are already stated in [8, Proposition 4.3]. In the lemma that follows, some properties of the twisted antipode are established.

Lemma 11. The following equalities hold:



Proof. Let us first show the relation $(a)_H$ Indeed, by definition of \tilde{S} , the anti-comultiplicativity of the antipod $\underline{PSrtherquassonia}$ tivity, and the naturality of the braiding, we have

$$\underbrace{\operatorname{PSfrag replacements}}_{H \ c} \xrightarrow{H}_{H \ c} \xrightarrow{H}_{H$$

Let us show the relation (b). Indeed, by definition of S, the fact that comultiplication is an algebra morphism, the fact that δ_H is an algebra morphism, and the naturality of the braiding we have replacements $\sim \delta \delta$

$$\frac{\text{PSfrag replacements}}{H} \bigoplus_{H} \bigoplus_{H}$$

Let us show the relation (c). Indeed, this relation follows by the definition of \tilde{S} , the coassociativity, the definition of \tilde{S} , the counterline \tilde{S} is the definition of \tilde{S} , the counterline \tilde{S} is the definition of \tilde{S} , the counterline \tilde{S} is the definition of \tilde{S} is the definition

Now we show the equality (d). It follows the part (a), the naturality of the braiding, the definition of \tilde{S} , the formula of \tilde{S} is an algebra morphism, and the definition of left coadjoint coaction:

$$\frac{\text{PSfrag replacements}}{H} \bigoplus_{i=1}^{d} \bigoplus_{i=1}^{d}$$

The equality (e) is a consequence of the equality (d). To see this, compose the left hand side of (d) with the antipode S of H and use the definition of \tilde{S} . HFinally, let us show the equation (f). Indeed, this equation follows by the part (b), the fact that comultiplication is an algebra morphism, the (co) associativity, the naturality of



Remark 12. Another useful property of the twisted antipode \tilde{S} is that $\varepsilon \tilde{S} = \delta$. It follows by the definition of \tilde{S} , the fact that $\varepsilon S = \varepsilon$, and the counitality.

The following lemma gives the expression of the paracocyclic operator $\tau_n(\delta, \sigma)$ in terms of $\tau_{n-1}(\varepsilon, u)$.

Lemma 13. If $n \ge 2$, then

Proof. Let us first show the equation H^{a} . Indeed, by definition $H^{a}(\delta, \sigma)$, Lemma 11(*a*), the naturality of the braiding, inductive definition of the left diagonal action, and the definition of $S_{THE}^{a}(\varepsilon, u)$, we have $\sim \sim \sim \sigma_{H}^{a}$

$$\frac{PSfrag replacements}{\tau_n(\delta,\sigma)} = \begin{pmatrix} H^{\otimes n-1} & H & H^{\otimes n-1} & H^{\otimes n-1} \\ H^{\otimes n-1} & H^{\otimes n-1} & H^{\otimes n-1} & H^{\otimes n-1} \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} & \sigma \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} & \sigma & H^{\otimes n-2} \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} & \sigma & H^{\otimes n-2} \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} & \sigma & H^{\otimes n-2} \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} & \sigma & H^{\otimes n-2} \\ H^{\otimes n-1} & \sigma & H^{\otimes n-1} \\ T_{n-1}(\varepsilon, u) & T_{n-1}(\varepsilon, u) \\ T_{n-1}(\varepsilon, u) & T_{$$

Further, we show the part (b). For n = 2, the statement follows by definition. From now on, suppose the strategy the definition of $\pi_n(\delta, \sigma)$, the definition of the left diagonal action, the coassociativity, and the definition of $\tau_{n-1}(\varepsilon, u)$, we have:



The equalities stated in the following lemma are used in computation of squares of the paracocyclic operator $\tau_n(\delta, \sigma)$ in the case $n \geq 3$.





Proof. We begin by showing the equality (a). Let us first inspect the case n = 2. To see that the equality is true in this case, $\frac{PStrag}{rest}$ is the case n = 2. To comultiplication is an algebra single plasments the coassociate view of $\tau_{2}(\varepsilon, u)$, the fact that the antipode, and the naturality of the braiding: $\tau_{2}(\varepsilon, u)$

From now on, suppose that $\frac{PSf_{rag}}{T} = \frac{PSf_{rag}}{T} = \frac{$



Let us show the equality (b). Indeed, it follows by definition of $\frac{1}{1}n(\varepsilon, u)$, the naturality of the braiding and the associativity: n-1





Here (i) follows by definition of $\tau_n(\varepsilon, u)$ and $\tau_{n-1}(\varepsilon, u)$, (ii) follows by inductive definition of left diagonal action, (iii) follows by the anti-multiplicativity of the antipode, (iv) follows by the fact that multiplication is an algebra morphism, (v) follows by the axiom of a module and the anti-comultiplicativity of the antipode, (vi) follows by the naturality of the braiding, the (co)associativity, and by the axiom of a module, (vii) follows by the fact that $\varepsilon S = \varepsilon$, the naturality of the braiding, and definition of $\tau_{n-1}(\varepsilon, u)$.

The equalities from the following lemma show how the endomorphism $m(\mathrm{id}_H \otimes \sigma)$ interacts with the paracocyclic operator $\tau_n(\varepsilon, u)$. These equalities are intensively used while proving Formula (5) by using Formula (4) of Theorem 2.





Here (i) follows by definition of $\tau_n(\varepsilon, u)$, (ii) follows by the anti-multiplicativity of antipode, (iii) follows by the naturality of the braiding and the fact that multiplication is an algebra **memphism** (iii) follows by the anti- $\overset{n}{\mathcal{C}}$ multiplicativity of $\overset{n}{\sigma}$ ntipode, (v) follows by the fact that σ is a coalgebra morphism and by the axiom of a module and (vi) follows by definition of $\tau_n(\varepsilon, u)$ and the naturality $\breve{\sigma}$ (the braiding. $\frac{2}{i}$

Let us now show the part (b). Indeed, by definition of $\tau_n(\varepsilon, u_n^{j})$ and the left diagonal <u>PSfrag replacetives</u>, by the naturality of the braiding, and the associativity, we have:



Before passing to the proof of Theorem 2, let us state another auxiliary lemma.

Lemma 16. We have the following assertions:



The equality (b) from Lemma 16 is intensively used while proving both of the equalities from Theorem 2. The equality (c) from Lemma 16 is particularly used in final steps of the computation of $\tau_n(\delta, \sigma)^{n+1}$.

Proof. Let us first show the part (a). We first show that the morphism from part (a) is an algebra morphism. By using definition of the left coadjoint coaction, the bialgebra



Similarly, by using definition of the left coadjoint coaction, the fact that unit is a coalgebra morphism, the fact that δ is an algebra **Examplify** due to the fact that Su = u, we have:



Let us now show that the morphism from (a) is a $\frac{\text{PSfrag replacements}}{\text{coalgebra morp}}$ hism. Indeed, by definition of the left coadjoint coaction, the fact that coalses are algebra morphism, the naturality of the braiding, the coassociativity the antipode axiom, and the (co) unitality, we have:

Furthermore, by definition of the left coal straight endowed with the fact that δ is an algebra morphism, the naturality of the parading the (co)unitality, and the antipode axiom we have: have: $\delta = \frac{1}{\delta} \frac{H}{\delta}$

$$\frac{PSfrag replacements}{H} \underbrace{\partial \left(\begin{array}{c} \delta \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\begin{array}{c} A \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\left(\begin{array}{c} A \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\left(\begin{array}{c} A \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\left(\begin{array}{c} A \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\left(\begin{array}{c} A \\ H \end{array} \right)}_{H} \underbrace{\partial \left(\left(\begin{array}{c} A \\$$

This completes the proof of the part (a).

Let us show the part (b) by induction. For n = 1, we prove the statement as follows. By using the part (a), the definition of the coadjoind statements fact that δ is an algebra morphism, the naturality of the braiding the coasso statements fact that δ is an algebra δ (co)unitality we have: H H H



Suppose that the statement is true for an $n \ge 1$ and let us show it for n+1. We have:





which shows the desired statement. Here (i) and (viii) both follow by inductive definition of left diagonal action and left coadjoint coaction, (ii) follows by the fact that δ is an algebra morphism, (iii) follows by the coassociativity, (iv) follows by the naturality of the braiding and the case n = 1, (v) and (vii) both follow by the naturality of the braiding and by the coassociativity and (vi) follows by the induction hypothesis.

Finally, we show the part (c) by induction. For n = 1, the statement follows by definition of right adjoint action and the fact that σ is a coalgebra <u>PhirphremeStrippose</u> that the statement is true for $n \geq 1$ and let up<u>shagweptacements</u> + 1. Indeed, $\overset{H\otimes n+1}{\text{by}}$ using inductive definition of the left and the right diagonal actions, the anti-multiplicativity of the antipode, the fact that σ is a coalgebra morphism, the naturality of the braiding, for the induction hypothesis, we have: $\overset{H}{\text{potential}}$ $\overset{H}{\text{potential}}$ $\overset{H}{\text{potential}}$ $\overset{H}{\text{potential}}$



Compare 17. If (o, σ) is a modular pair, then $\stackrel{\text{PSfrag replacements}}{\overset{\text{O}}{\text{o}}} \stackrel{\text{TH}}{\overset{\text{O}}{\text{o}}} \stackrel{\text{Indeed, this follows by the}}{\overset{\text{PSfrag replacements}}{\overset{\text{O}}{\text{o}}} \stackrel{\text{TH}}{\overset{\text{O}}{\text{o}}} \stackrel{\text{Indeed, this follows by the}}{\overset{\text{PSfrag replacements}}{\overset{\text{PSfrag replacements}}}}$ by the definition of \tilde{S} , the fact that σ is a goalgebra/morphism, and since the (δ, σ) is $\stackrel{\text{H}}{\overset{\text{H}}{\text{o}}}$ modular pair: $\stackrel{\text{PSfrag replacements}}{\overset{\text{O}}{\text{H}} \stackrel{\sigma}{\overset{\text{O}}{\text{o}}} \stackrel{\text{H}}{\overset{\text{O}}{\text{H}}} \stackrel{\sigma}{\overset{\text{O}}{\text{H}}} \stackrel{\sigma}{\overset{\sigma}{\overset{\text{O}}{\text{H}}}} \stackrel{\sigma}{\overset{\sigma}{\overset{\sigma}{\overset{\sigma}}} \stackrel{\sigma}{\overset{\sigma}{\overset{\sigma}{\overset{\sigma}}}}$

$$\begin{array}{c} H \textcircled{\bigcirc} = H \textcircled{\bigcirc} \textcircled{\bigcirc} = H \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} = H \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} P \underline{\operatorname{Sfr}} \underbrace{\operatorname{\$geplacements}}_{H} \\ \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \textcircled{\bigcirc} \overset{H} \textcircled{\bigcirc} \overset{H} \textcircled{\bigcirc} \overset{H} \underset{H}{\overset{H}} \end{array}$$

Η

6.2. **Proof of Formula** (4). The proof of Formula (4) of Theorem 2 is divided into several steps. For n = k = 2, it suffices to calculate the square of $\tau_2(\delta_{\mathcal{H}}^{\sigma})$. For $n \geq_{\mathcal{H}}^{H}$, we first calculate the square and then derive formulas for the remaining powers.



Here (i) follows by definition of $\tau_2(\delta, \sigma)$, (ii) follows from Lemma 13(a) for n = 2 and since $\tau_1(\varepsilon, u) = S$, (iii) follows by the fact that multiplication is an algebra morphism

	PSfrag replacements
PSfrag replacements	$\tau_{n-1}(\varepsilon, u)_1$
1	12
n-1	<i>n</i> –21

ON THE BRAIDED CONNES-MOSCOVICI CONSTRUCTION

and associativity, (iv) by the anti-multiplicativity of the antipode, (v) follows by the associativity and the naturality of the braiding and (vi) follows by the antipode axiom, the naturality of the braiding, the (co)finitality, and the fact that $f_{i}^{*}(\varepsilon, u) = S$.

From now on, let us assume that $\tau_{\tau_{\tau_n}} = 0$. PSfrag replacements δ_1 . Let us calculate the sequence of $\tau_n(\delta, \sigma)$. We have $\tau_{\pi_n} \downarrow (\xi \in \psi)$



which indeed shows Formula (4) in the case $n \ge 3$ and k = 2. Here $\frac{1}{2}(i)$ follows by parts (a) and (b) of Lemma 13, (ii) follows by the associativity and from Lemma 14(a) for n-1, (iii) follows by the associativity and from Lemma 14(b) for n-1, (iv) follows from Lemma 14(c) for n-1, (v) follows from Lemma 14(d) for n-1 and (vi) follows_{σ} by the counitality, definition of the twisted antipode \tilde{S} , and from Lemma 13(b) applied on $\delta = \varepsilon$ and $\sigma = u$ for n-1.

$$\frac{\text{PSfrag replacements}}{1} \qquad \qquad \tau_{n-1}(\varepsilon, u)$$

6.2.2. Passing from $(\tau_n(\delta, \sigma))^2$ to $(\tau_n(\delta, \sigma))^3$, $n \ge 3$. From the calculation that has been done in Section 6.2.1, we can easily deduce Formula (4) for $n \ge 3$ and k = 3.





Here (i) follows by developing $\tau_n(\delta, \sigma)^3$ as $\tau_n(\delta, \sigma)^2 \tau_n(\delta, \sigma)$, by definition of $\tau_n(\delta, \sigma)$ and by the computation of $\tau_n(\delta, \sigma)^2$ that has Been determine 2.1, (ii) follows by the coassociativity and Lemma 11(f), (iii) follows by the naturality of the braiding and the coassociativity, (iv) follows from Lemma 15(b) for j = n-1 and from Lemma 11(d), (v) follows by the naturality of the braiding and the definition of $\tau_{n-1}(\varepsilon, u)$.

6.2.3. Computation of $(\tau_n(\delta, \sigma))^{j+1}$, $2 \leq j \leq n - 1$. Note that by now, we have completely shown Formula (4) in the cases n = 2 and n = 3. Also, the square and the cube of $\tau_n(\delta, \sigma)$ are calculated for each $n \geq 3$. In this section, we finish the proof of (4), by focusing on the case $n \geq 4$. As it has been already noted, for $j_{H^{\otimes n-2}}$, $(\tau_n(\delta, \sigma))^{j+1}$ is already computed in Section 6.2.2. From now on, we assume that $3 \leq j \leq n - 1$. If (4) is established for k = j, then: $(\tau_{n-1}(\varepsilon, u))^{j-1}$





ON THE BRAIDED CONNES-MOSGOVICI CONSTRUCTION



which finishes the proof of (4). Here (i) follows by decomposing $(\tau_n(\delta,\sigma))^{j+1}$ in the composition $(\tau_n(\delta,\sigma))^j \tau_n(\delta,\sigma)$, by definition of $\tau_n(\delta,\sigma)$ and by the hypothesis that (4) is established for k = j, (ii) follows by the coassociativity and Lemma 11(f), (iii) follows by the naturality of the braiding and the coassociativity, (iv) follows from Lemma 16(b) and by applying j-1 times Lemma 15(b), (v) follows from Lemma 11(d) and the coassociativity and finally, (vi) follows by the naturality of the braiding and the definition of $\tau_{n-1}(\varepsilon, u)$.

6.3. **Proof of Formula** (5). In order to show Formula (5) of Theorem 2, we will separately consider three cases: n = 1, n = 2, and n > 3. PSfrag replacements

If n = 1, then we have



Here (i) follows by definition $\tau_1(\delta, \sigma)$, (ii) follows from Lemma 11(b), (iii) follows by the naturality of the braiding, the associativity, and Rem $\frac{17}{10}$ (iv) follows by definition of right adjoint action, the fact that σ is a coalgebra morphism, and Lemma 11(e).



Here (i) follows by expanding $(\tau_2(\delta, \sigma))^3$ as $(\tau_2(\delta, \sigma))^2 \tau_2(\delta, \sigma)$, using the definition of $\tau_2(\delta, \sigma)$, and Formula (4) in the case n = k = 2, (ii) follows from Lemma 11(f), (iii) by using the fact that σ is a coalgebra morphism, (iv) by the naturality of the braiding, definition

$$\frac{\text{PSfrag replacements}}{1}$$
34 IVAN BARTULOV $(\hat{\mathbf{f}}_{n-1}(\varepsilon, u))^{n-1}$

of $\tau_1(\delta, \sigma)$, and the anti-multiplicativity of the antipode, $\stackrel{\tau_{n-1}(\varepsilon, u)}{(v)}$ follows by the naturality of the braiding, by definition of right adjoint action, and the case n = 1 of Formula (5), which is shown above.

From now on, let us assume that $n \geq 3$. We have



which shows Formula (5). Here (i) follows by decomposing $(\tau_n(\delta, \sigma))^{n+1}$ in the composition $(\tau_n(\delta, \sigma))^n \tau_n(\delta, \sigma)$, by using Lemma 13(b), and Formula (4) for k = n, (ii) follows from Lemma 11(f), (iii) by the fact that σ is a coalgebra morphism, (iv) by definition of $\tau_{n-1}(\varepsilon, u)$, the naturality of the braiding, definition of $\tau_1(\delta, \sigma)$, and the coassociativity, (v) by definition of the twisted antipode, Lemma 16(b), and by applying n - 2 times Lemma 15(b), (vi) follows by the naturality of the braiding, by combining Lemma 15(a) with Lemma 16(c) for n - 1, by the coassociativity, and Lemma 11(d), (vii) follows by the naturality of the braiding and definition of $\tau_{n-1}(\varepsilon, u)$.

7. Proof of Theorem 7

In order to show that the family $\{\alpha_n \colon H^{\otimes n} \to C^{\otimes n+1}\}_{n \in \mathbb{N}}$ is a morphism between the paracocyclic objects $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ and $\mathbf{C}_{\bullet}(C)$ in \mathcal{B} , we will directly check that

(7)
$$\alpha_n \delta_i^n(\sigma) = \delta_i^n \alpha_{n-1} \text{ for } 0 \le i \le n, n \ge 1$$

(8)
$$\alpha_n \sigma_i^n = \sigma_i^n \alpha_{n+1} \quad \text{for} \quad 0 \le j \le n, n \ge 0,$$

(9)
$$\alpha_n \tau_n(\delta, \sigma) = \tau_n \alpha_n \quad \text{for} \quad n \ge 0.$$

Note that we abusively use the same notation for cofaces, codegeneracies, and paracocyclic operators of two different constructions. These should be understood from context. Roughly described, the equalities (7) and (8) follow by the fact that C is a coalgebra in the category of right H-modules. In order to show the equality (9) in the case $n \ge 2$, we will need the following computation: FSmag replacements



Proof. We prove the claim by induction. Let us first show it for n = 2. Indeed, by the right module axiom, the naturality of braiding and the fact that the comultiplication $\Delta_C \colon C \to C \otimes C$ is an *H*-linear morphism, we have $\Delta_{\text{Sfrag replacements}}$



Suppose that the claim is true for an $n \ge 2$ and the us show it for n+1. We have



which indeed proves the claim for n + 1. Here (i) follows by the inductive definition of the left diagonal action, (ii) follows by the induction hypothesis, (iii) follows by the right module axiom and (iv) follows by the naturality of the braiding and the fact that the comultiplication $\Delta_C \colon C \to C \otimes C$ is an *H*-linear morphism. \Box

7.1. **Proof of the equality** (7). Let us show the equality (7). If n = 1 and i = 0, then the equality (7) writes as $\alpha_1 \delta_0^1 = \delta_0^1 \alpha_0$, which follows by definitions and the right module axiom. Let n = i = 1. In this case, the equality (7) writes as $\alpha_1 \delta_1^1 = \delta_1^1 \alpha_0$, which is exactly the condition that $\alpha \colon \mathbb{1} \to C$ is a σ -trace. This shows the equality (7) for n = 1and $0 \leq i \leq 1$. PSfrag replacements

Now let $n \ge 2$ and i = 0. By definitions, the naturality of the braiding, the axiom of a right module, and the coassociativity, we have: Cn-1



Let $n \geq 2$ and $1 \leq i \leq n-1$. In this case, equation (7) follows by definitions, the coassociativity, and the fact that $\Delta_C: C \to C \otimes \mathbb{C}$ is an *H*-linear morphism:



Finally, let $n \ge 2$ and i = n. Equation (7) in this case follows by the case $n \ge 2$ and i = 0, which is written above, the equation (9), which is proven in Section 7.3, and by the paracyclic compatibility relation $\tau_n \delta_0^n = \delta_n^n$ (see Section 3.2). Indeed, we have

$$\alpha_n \delta_n^n = \alpha_n (\tau_n(\delta, \sigma) \delta_0^n) = (\alpha_n \tau_n(\delta, \sigma)) \delta_0^n = (\tau_n \alpha_n) \delta_0^n =$$
$$= \tau_n (\alpha_n \delta_0^n) = \tau_n (\delta_0^n \alpha_{n-1}) = (\tau_n \delta_0^n) \alpha_{n-1} = \delta_n^n \alpha_{n-1}.$$

7.2. **Proof of the equality** (8). Let $\underline{\text{usfshowplithereact}}$ uality (8). We consider the three following cases: $j = 0, 1 \leq j \leq n-1$, and j = n. In each case, the desired equality follows by definition, the counitality, the fact that the counit $\varepsilon_{R}^{C}: C \to \mathbb{1}$ is an *H*-linear morphism, and the naturality of the braiding.





37

7.3. **Proof of the equality** (9). Let us verify that equation (9) holds. When n = 0, this holds since twist morphisms are natural, $\theta_1 = \mathrm{id}_{1,H}$ and since $\tau_0(\delta, \sigma)_{H} = \mathrm{id}_{1}$. Indeed, $\frac{\mathrm{PSfrag \ replacements}}{\mathrm{PSfrag \ replacements}} = \theta \quad \alpha = \alpha = \delta \quad \tau \quad (\delta, \sigma) = \delta$

$$\frac{\frac{eplacements}{\tau_0 \alpha_0}}{C} = \theta_C \alpha_0 = \alpha_0 = \overset{A}{\underset{C}{\otimes}} \tau_0(\delta, \sigma).$$

C

 α

Let us check it for the case $n = 1\alpha$ Indeseds we have the set of the case $n = 1\alpha$ indeseds we have the set of the set



Here (i) and (viii) follow by definition, (ii) follows by the fact that α is δ -invariant, (iii) follows from the fact that the comultiplication $\Delta_C: C \to C \otimes C$ is *H*-linear. The equality (iv) follows by the (co)associativity, (v) follows by the antipode axiom and the (co)unitality, (vi) follows from the fact that α is a σ -trace and (vii) follows by the naturality of the braiding.



Here (i) and (x) follow by definition, (ii) follows by applying Lemma 18, (iii) follows from the fact that α is δ -invariant, (iv) follows from the fact that the comultiplication $\Delta_C: C \to C \otimes C$ is H-linear. The equality (v) follows from the coassociativity and the right module axiom, (vi) follows by the antipode axiom, the (co)unitality, and the right module axiom, (vii) follows by the coassociativity and the naturality of the braiding, (viii) follows from the fact that α is a σ -trace and finally, (ix) follows by the naturality of the braiding.

8. Appendix

In this appendix, we verify paracyclic compatibility relations (PCR) of the paracocyclic object $\mathbf{CM}_{\bullet}(H, \delta, \sigma)$ defined in Section 4.3. Note that the verification of simplicial relations (SR) for this object is an easy task. One can show it graphically, by using the level-exchange property (see Section 2.4), the coassociativity, and the counitality. Also, the relation $\tau_n \sigma_i^n = \sigma_{i-1}^n \tau_{n+1}$ for $1 \leq i \leq n$ follows from the bialgebra axiom and by the naturality of the braiding. In this appendix, we show that

- (1) $\tau_n \delta_i^n = \delta_{i-1}^n \tau_{n-1}$ for $n \in \mathbb{N}^*$ and $1 \le i \le n$, (2) $\tau_n \delta_0^n = \delta_n^n$ for $n \in \mathbb{N}^*$, (3) $\tau_n(\delta, \sigma) \sigma_0^n = \sigma_n^n (\tau_{n+1}(\delta, \sigma))^2$ for $n \in \mathbb{N}$.

Verifications of (1)-(3) are somewhat involved. Note that (3) is proved for n = 2 in [8, Theorem 7.1.]. Recall the notion of twisted antipodes and the corresponding notation from Section 6.

8.1. The relation $\tau_n \delta_i^n = \delta_{i-1}^n \tau_{n-1}$. If n = 1 and i = 1, the relation rewrites as $\tau_1 \delta_1^1 = \delta_0^1 \tau_0$ and it follows by definitions, the fact that σ is a coalgebra morphism, the fact that (δ, σ) is a modular pair and by the antipode axiom:



If n = 2 and i = 1, the relation rewrites as $\tau_2 \delta_1^2 = \delta_0^2 \tau_1$ and it follows by definitions, Lemma 11*a*), the co**responsibility** entropy antipode axiom, and the naturality of the braiding:



If n = 2 and i = 2, the relation rewrites as $\tau_2 \sigma_2^{-1} = \sigma_1^{-1} \frac{1}{H} \frac{1}{H}$ by definitions, the fact that σ is a coalgebra morphism, and the fact that multiplication $H_{\mathcal{H}^{\otimes n-2}}$ an algebra morphism: PSfrag replacements

$$\frac{PSfrag replacements}{1} \xrightarrow{H} \xrightarrow{PSfrag replacements} H \xrightarrow{PSfrag repl$$

Now let $n \geq 3$. For i = 1, the relation rewriting as $\tau_n \delta_1^n = \delta_0^n \tau_{n-1}^{H^{\otimes n-1}}$ and it is true since



Here (i) and (ix) follow by definition, (ii) follows by inductive definition of the left diagonal action, (iii) and (viii) follow from Lemma 11(a), (iv) and (vi) follow by the naturality of the braiding and the coassociativity, (v) follows by the anti-comultiplicativity of the antipode, (vii) follow by the antipode axiom, the counitality, and the naturality of the braiding. For $2 \leq i \leq n-1$, the relation $\tau_n \delta_i^n = \delta_{i-1}^n \tau_{n-1}$ is a consequence of the fact that multiplication is an algebra morphism and the naturality of the braiding. Let us check $\tau_n \delta_i^n = \delta_{i-1}^n \tau_{n-1}$ for i = n. The relation follows by definition, Lemma 13b), by the fact that σ is a coalgebra

$$\begin{array}{c} \underline{\operatorname{PSfrag\ replacements}}\\ H\\ H^{\otimes n-2}\\ \delta\\ \\ \underline{\operatorname{PSfrag\ replacements}} \text{IVAN\ BARTELQVL}\\ H\\ \\ \end{array}$$
rphism, and by the fact that Hend complete comultiplication is an algebra morphism:
$$\begin{array}{c} \delta\\ \delta\\ \delta\\ \delta\\ \end{array}$$

40

mo

$$\frac{\text{PSfrag replacements}}{\tau_n \delta_n^n} = \underbrace{\left\{ \begin{array}{c} H^{\otimes n-2} \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} H^{\otimes n-2} \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ \hline \tau_{n-1}(\varepsilon, u) \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ \sigma \\ H \end{array}\right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ H \end{array} \right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\ H \end{array}\right\}}_{H H^{\otimes n-2}} \underbrace{\left\{ \begin{array}{c} \sigma \\$$

8.2. The relation $\tau_n \delta_0^n = \delta_n^n$. For n = 1, the relation rewrites as $\tau_1 \delta_0^1 = \delta_1^1$ and it follows by definition, the fact that δ is an algebra morphism, by $\varepsilon S = \varepsilon$, and the fact that the unit is a coalgebra morphism:



Now let $n \ge 2$. By definition, the fact that δ is an algebra morphism, by $\varepsilon S = \varepsilon$, the fact that the unit is a coalgebra morphism, the idente-module axiom, and the naturality of the braiding, we have $\sum_{\text{Sfrag replacements}} \frac{n-1}{\delta} = \frac{n-1}{\delta}$



8.3. The relation $\tau_n(\delta, \sigma)\sigma_0^n = \sigma_n^n(\tau_{n+1}(\delta, \sigma))_{\delta}^{\underline{H}}$. In order to show the relation, one can use Theorem 2. We first prove the case n = 0. It is true since δ



Here (i) follows by using Formula (5) of Theorem 2 and the definition of σ_0^0 , (ii) follows from definition of right adjoint action and left coadjoint action of H on itself, the naturality of the braiding, and the fact that δ is an algebra morphism and σ is a coalgebra morphism. The equality (iii) follows by applying twice the fact that the counit is an algebra morphism, (iv) follows by the fact that $\varepsilon S = \varepsilon$ and since σ is a coalgebra morphism, (v) follows by the fact that δ is an algebra morphism and by the antipode axiom, (vi) follows by definition of $\tau_0(\delta, \sigma)$ and σ_0^0 .



Here (i) follows by Theorem 2 applied for n + 1 and k = 2 and by definition of σ_n^n . The equality (ii) follows by the fact that counit is an algebra morphism and the naturality of the braiding, (iii) follows from Lemma 11(b) and the fact that σ is a coalgebra morphism, (iv) follows from Lemma 11(e), by the fact that counit is an algebra morphism, and by the naturality of the braiding. The equality (v) follows by the fact that $\varepsilon S = \varepsilon$, by Remark 12, and Lemma 16(a), (vi) follows by definition of the paracocyclic operator $\tau_n(\delta, \sigma)$ and finally, (vii) follows by definition of $\tau_n(\delta, \sigma)$ and σ_0^n .

References

- S. E. Akrami and S. Majid. Braided cyclic cocycles and nonassociative geometry. Journal of Mathematical Physics, 45(10):3883–3911, 2004.
- [2] A. M. Bohmann, T. Gerhardt, A. Hogenhaven, B. Shipley, and S. Ziegenhagen. Computational tools for topological coHochschild homology. *Topology and its Applications*, 235:185–213, 2018.
- [3] A. Connes. Cohomologie cyclique et foncteurs Ext. 1983.
- [4] A. Connes. Non-commutative differential geometry. Publications Mathématiques de l'Institut des Hautes Études Scientifiques, 62(1):41–144, Dec. 1985.
- [5] A. Connes and H. Moscovici. Hopf Algebras, Cyclic Cohomology and the Transverse Index Theorem. Communications in Mathematical Physics, 198(1):199–246, Nov. 1998.
- [6] A. Connes and H. Moscovici. Cyclic Cohomology and Hopf Algebras. Letters in Mathematical Physics, 48(1):97–108, Apr. 1999.
- M. Farinati and A. Solotar. Cyclic cohomology of coalgebras, coderivations and De Rham cohomology. In *Hopf algebras and quantum groups*, volume 209 of *Lect. Notes in Pure and Applied Math.*, pages 105–130. Marcel Dekker, 2000.

IVAN BARTULOVIĆ

- [8] M. Khalkhali and A. Pourkia. Hopf cyclic cohomology in braided monoidal categories. Homology, Homotopy and Applications, 12(1):111 – 155, 2010.
- [9] S. M. Lane. Natural associativity and commutativity. *Rice Institute Pamphlet-Rice University Studies*, 49(4), 1963.
- [10] S. M. Lane. Homology. Springer, 1995.
- B. L. Tsygan. The homology of matrix Lie algebras over rings and the Hochschild homology. *Russian Mathematical Surveys*, 38(2):198, 1983.
- [12] V. Turaev and A. Virelizier. Monoidal Categories and Topological Field Theory. Birkhäuser, 2017.
- [13] C. A. Weibel. An introduction to homological algebra. Number 38. Cambridge University Press, 1995.

UNIVERSITÉ DE LILLE, LABORATOIRE PAUL PAINLEVÉ UMR CNRS 8524, F-59000 LILLE, FRANCE *Email address*: ivan.bartulovic@univ-lille.fr