FUNCTIONAL A POSTERIORI ERROR ESTIMATES FOR ELLIPTIC PROBLEMS IN EXTERIOR DOMAINS

Dirk Pauly and Sergey Repin

November 21, 2018

Abstract

This paper is concerned with the derivation of computable and guaranteed upper bounds of the difference between the exact and the approximate solution of an exterior domain boundary value problem for a linear elliptic equation. Our analysis is based upon purely functional argumentation and does not attract specific properties of an approximation method. Therefore, the estimates derived in the paper at hand are applicable to any approximate solution that belongs to the corresponding energy space. Such estimates (also called error majorants of the functional type) have been derived earlier for problems in bounded domains of \mathbb{R}^N (see [2, 3]).

Key Words A posteriori error estimates of functional type, elliptic boundary value problems in exterior domains

AMS MSC-Classifications 65 N 15

Contents

1	Introduction	2
2	Upper bounds for the deviation from the exact solution in dimensions $N > 2$	5
	$N \geq 3$	
	2.1 First estimate	
	2.2 Second estimate	7
	2.3 Third estimate	8
3	Upper bounds in dimension $N=2$	10
\mathbf{A}	Appendix	10
	A.1 Lower bounds for the error	10
	A 2 Poincare type estimates for exterior domains	11




Figure 1: exterior domain Ω with artificial interface Γ

1 Introduction

The main focus of our investigations is to suggest a method of deriving guaranteed and computable upper bounds of the difference between the exact solution u of an elliptic exterior domain boundary value problem and any approximation from the corresponding energy space. We discuss the method with the paradigm of the prototypical elliptic problem

$$-\operatorname{div} A\nabla u = f \qquad \qquad \text{in } \Omega, \tag{1.1}$$

$$u|_{\gamma} = g$$
 on $\gamma := \partial \Omega$. (1.2)

We assume that $\Omega \subset \mathbb{R}^N$ with $N \geq 1$ is an exterior domain, i.e. $\mathbb{R}^N \setminus \Omega$ is compact, with Lipschitz continuous boundary γ (see Figure 1).

Throughout this paper we will use the weighted Lebesgue function spaces

$$\mathsf{L}^2_s(\Omega) := \left\{ \varphi \ | \ \rho^s \varphi \in \mathsf{L}^2(\Omega) \right\}, \quad s \in \mathbb{R}.$$

Here $\rho:=(1+r^2)^{1/2}$ and r(x):=|x| denotes the radius vector. $\mathsf{L}^2_s(\Omega)$ is a Hilbert space equipped with the scalar product

$$\langle \varphi, \psi \rangle_{s,\Omega} := \langle \rho^s \varphi, \rho^s \psi \rangle_{\Omega} := \int_{\Omega} \rho^{2s} \varphi \psi \, \mathrm{d}\lambda,$$

where φ and ψ belong to $\mathsf{L}^2_s(\Omega)$ and λ is Lebesgue's measure. We denote the corresponding norms by $\|\varphi\|_{s,\Omega} = \|\rho^s\varphi\|_{\Omega}$. If s=0 then $\mathsf{L}^2_s(\Omega)$ coincides with the usual Lebesgue space

 $L^2(\Omega)$. For the sake of simplicity we keep the same notation for spaces of vector-valued functions. Moreover, we introduce the weighted Sobolev space

$$\mathsf{H}^1_{-1}(\Omega) := \left\{ \varphi \in \mathsf{L}^2_{-1}(\Omega) \ | \ \nabla \varphi \in \mathsf{L}^2(\Omega) \right\},\,$$

which is a Hilbert space as well with respect to the scalar product

$$(\varphi, \psi) \mapsto \langle \varphi, \psi \rangle_{-1,\Omega} + \langle \nabla \varphi, \nabla \psi \rangle_{\Omega}$$

By $\mathring{\mathsf{H}}^1_{-1}(\Omega)$ we denote the closure of $\mathring{\mathsf{C}}^\infty(\Omega)$, the space of compactly supported smooth test functions, in the norm of $\mathsf{H}^1_{-1}(\Omega)$. Whenever we consider Sobolev spaces of bounded domains we use the usual unweighted L^2 -scalar products and -norms.

For dimensions $N \geq 3$ the solution theory for the problem (1.1)-(1.2) is based on the weighted Poincare/Friedrich estimate (see Corollary 16 (i) and Remark 17 of the appendix)

$$\|\varphi\|_{-1,\Omega} \le \frac{2}{N-2} \|\nabla\varphi\|_{\Omega} \quad \forall \, \varphi \in \overset{\circ}{\mathsf{H}}_{-1}^{1}(\Omega), \tag{1.3}$$

the Lax-Milgram theorem and, if needed, an adequate extension operator for the boundary data. Let u_{γ} be some function in $\mathsf{H}^1_{-1}(\Omega)$ satisfying the boundary condition (1.2). The weak solution $u \in \overset{\circ}{\mathsf{H}}^1_{-1}(\Omega) + u_{\gamma} \subset \mathsf{H}^1_{-1}(\Omega)$ of (1.1)-(1.2) is then defined by the variational formulation

$$\langle A\nabla u, \nabla w \rangle_{\Omega} = \langle f, w \rangle_{\Omega} \quad \forall w \in \overset{\circ}{\mathsf{H}}_{-1}^{1}(\Omega).$$
 (1.4)

By (1.3) the left hand side of (1.4) is a strongly coercitive sesqui-linear form over $H_{-1}^1(\Omega)$ provided that the real-matrix-valued function A is measurable, bounded a.e., symmetric and uniformly strongly elliptic, i.e.

$$\exists c_A > 0 \quad \forall \xi \in \mathbb{R}^N \quad \forall x \in \Omega \quad c_A |\xi|^2 \le A(x)\xi \cdot \xi. \tag{1.5}$$

If $f \in \mathsf{L}^2_1(\Omega)$ then by the Cauchy-Scharz inequality the right hand side of (1.4) is a linear and continuous functional over $\mathsf{H}^1_{-1}(\Omega)$. Thus, under these assumptions the problem (1.4) is uniquely solvable in $\mathsf{H}^1_{-1}(\Omega) + u_{\gamma}$ by Lax-Milgram's theorem.

If N=1,2 one can apply the same arguments with the difference that (1.3) has to be modified. For N=1 and, for example, $\Omega\subset\mathbb{R}_+$ we have by Corollary 16 (iii) and Remark 17

$$\|\varphi\|_{-1,\Omega} \le 2 \, \|\varphi'\|_{\Omega} \quad \forall \, \varphi \in \overset{\circ}{\mathsf{H}}^{1}_{-1}(\Omega). \tag{1.6}$$

Hence, we get the same solution theory with tiny restrictions on Ω , which easily can be removed by a translation. For N=2 the singularities are stronger and additionally we

have to utilize logarithmic terms. By Corollary 16 (ii) and Remark 17 we have for domains $\Omega \subset \mathbb{R}^2$, such that the complement $\mathbb{R}^2 \setminus \Omega$ contains the unit ball,

$$\|\varphi/(r\ln r)\|_{\Omega} \le 2 \|\nabla\varphi\|_{\Omega} \quad \forall \varphi \in \overset{\circ}{\mathsf{H}}_{-1,\ln}^{1}(\Omega), \tag{1.7}$$

where

$$\mathsf{H}^1_{-1,\ln}(\Omega) := \left\{ \varphi \ | \ \varphi/(r\ln r), \nabla \varphi \in \mathsf{L}^2(\Omega) \right\}$$

is a Hilbert space equipped with the natural scalar product

$$(\varphi, \psi) \mapsto \langle \varphi/(r \ln r), \psi/(r \ln r) \rangle_{\Omega} + \langle \nabla \varphi, \nabla \psi \rangle_{\Omega}$$

and again $\overset{\circ}{\mathsf{H}}^1_{-1,\ln}(\Omega)$ denotes the closure of $\overset{\circ}{\mathsf{C}}^{\infty}(\Omega)$ in the norm of $\mathsf{H}^1_{-1,\ln}(\Omega)$. Consequently, we obtain for all f with $r \ln r f \in \mathsf{L}^2(\Omega)$ and all u_{γ} in $\mathsf{H}^1_{-1,\ln}(\Omega)$ satisfying the boundary condition (1.2) a unique solution u belonging to $\overset{\circ}{\mathsf{H}}^1_{-1,\ln}(\Omega) + u_{\gamma}$.

We summarize the results in the following

Theorem 1 Let $N \geq 3$ as well as $f \in \mathsf{L}^2_1(\Omega)$ and $u_{\gamma} \in \mathsf{H}^1_{-1}(\Omega)$ satisfying the boundary condition (1.2). Then the exterior boundary value problem (1.1)-(1.2) is uniquely weakly solvable in $\mathsf{H}^1_{-1}(\Omega) + u_{\gamma}$. The solution operator is continuous.

From the above discussion, it is clear that for N=1,2 the existence of weak solutions in suitable spaces can also be proved.

Remark 2 The boundary data g and its extension u_{γ} can be described in more detail. In the bounded domain case it is well known that there exists a bounded linear trace operator and a corresponding bounded linear extension operator (right inverse) mapping $H^1(\Omega)$ to $H^{1/2}(\gamma)$ and vice verse. Hence, by restriction we get a bounded linear trace operator

$$\tau_{\gamma}:\mathsf{H}^{1}_{-1}(\Omega)\to\mathsf{H}^{1/2}(\gamma)$$

and by extension and applying an obvious cutting technique we obtain a bounded linear extension operator

$$E: \mathsf{H}^{1/2}(\gamma) \to \mathsf{H}^1_{-1}(\Omega)$$

for our exterior domain Ω , which even maps to functions with (arbitrarily thin) compact support. As in the bounded domain case, E is a right inverse of τ_{γ} . Then we may specify $g \in \mathsf{H}^{1/2}(\gamma)$ and $u_{\gamma} := Eg \in \mathsf{H}^1_{-1}(\Omega)$ as well as our variational formulation for $u = \tilde{u} + Eg$: Find $\tilde{u} \in \overset{\circ}{\mathsf{H}}^1_{-1}(\Omega)$, such that

$$B(\tilde{u},w) := \langle A\nabla \tilde{u}, \nabla w \rangle_{\Omega} = \langle f, w \rangle_{\Omega} - \langle A\nabla Eg, \nabla w \rangle_{\Omega} =: F(w) \quad \forall \, w \in \overset{\circ}{\mathsf{H}}^{1}_{-1}(\Omega).$$

Finally, we introduce

$$\mathsf{D}(\Omega) := \left\{ \varphi \in \mathsf{L}^2(\Omega) \ | \ \operatorname{div} \varphi \in \mathsf{L}^2_1(\Omega) \right\},$$

which is a Hilbert space with respect to the canonical scalar product

$$(\varphi, \psi) \mapsto \langle \varphi, \psi \rangle_{\Omega} + \langle \operatorname{div} \varphi, \operatorname{div} \psi \rangle_{1,\Omega}$$
.

2 Upper bounds for the deviation from the exact solution in dimensions $N \geq 3$

Let v be an approximation of $u \in \overset{\circ}{\mathsf{H}}^1_{-1}(\Omega) + u_{\gamma} \subset \mathsf{H}^1_{-1}(\Omega)$, where v is assumed just to belong to $\mathsf{H}^1_{-1}(\Omega)$ since the boundary condition may not be satisfied exactly. Our goal is to obtain upper bounds for the difference between ∇u and ∇v in terms of the norm

$$\|\varphi\|_{A,\Omega} := \|A^{1/2}\varphi\|_{\Omega} = \langle A\varphi, \varphi \rangle_{\Omega}^{1/2}.$$

We use (1.4) and get for all $w \in \overset{\circ}{\mathsf{H}}_{-1}^1(\Omega)$

$$\langle A\nabla(u-v), \nabla w \rangle_{\Omega} = \langle f, w \rangle_{\Omega} - \langle A\nabla v, \nabla w \rangle_{\Omega}.$$
 (2.1)

Before we proceed we note two useful results.

Theorem 3 Let $u, v \in \mathsf{H}^1_{-1}(\Omega)$ be as above. Moreover, let Φ be a linear and continuous functional over $\overset{\circ}{\mathsf{H}}^1_{-1}(\Omega)$ and $c_{\Phi} > 0$, such that for all $w \in \overset{\circ}{\mathsf{H}}^1_{-1}(\Omega)$

$$\langle A\nabla(u-v), \nabla w \rangle_{\Omega} = \Phi(w) \le c_{\Phi} \|\nabla w\|_{A,\Omega}$$

holds. Then

$$\|\nabla(u - v)\|_{A,\Omega} \le c_{\Phi} + 2 \|\nabla(\hat{u} - \hat{v})\|_{A,\Omega} \tag{2.2}$$

for all $\hat{u}, \hat{v} \in \mathsf{H}^1_{-1}(\Omega)$, for which $\hat{u} - \hat{v}$ coincides with u - v on the boundary γ . If additionally u - v belongs to $\overset{\circ}{\mathsf{H}}^1_{-1}(\Omega)$ then

$$\|\nabla(u-v)\|_{A,\Omega} \le c_{\Phi}. \tag{2.3}$$

Proof We consider

$$w := u - v - (\hat{u} - \hat{v}) \in \overset{\circ}{\mathsf{H}}^{1}_{-1}(\Omega).$$

Using Cauchy-Schwarz' inequality we obtain

$$\|\nabla w\|_{A,\Omega}^{2} = \langle A\nabla(u-v), \nabla w \rangle_{\Omega} - \langle A\nabla(\hat{u}-\hat{v}), \nabla w \rangle_{\Omega}$$

$$\leq \left(c_{\Phi} + \|\nabla(\hat{u}-\hat{v})\|_{A,\Omega}\right) \|\nabla w\|_{A,\Omega}$$

and thus $\|\nabla w\|_{A,\Omega} \leq c_{\Phi} + \|\nabla(\hat{u} - \hat{v})\|_{A,\Omega}$. By the triangle inequality we get (2.2). (2.3) is trivial since we can set w := u - v, i.e. $\hat{u} := \hat{v} := 0$.

We may be more specific using the trace and extension operators from Remark 2.

Corollary 4 Let the assumptions of Theorem 3 be satisfied. Then

$$\|\nabla(u-v)\|_{A,\Omega} \le c_{\Phi} + 2 \|\nabla E(g-\tau_{\gamma}v)\|_{A,\Omega} \le c_{\Phi} + 2c_{\gamma} \|g-\tau_{\gamma}v\|_{\mathsf{H}^{1/2}(\gamma)}$$
.

Here $c_{\gamma} > 0$ is the constant in the inequality

$$\|\nabla E\varphi\|_{A,\Omega} \le c_{\gamma} \|\varphi\|_{\mathsf{H}^{1/2}(\gamma)} \quad \forall \varphi \in \mathsf{H}^{1/2}(\gamma). \tag{2.4}$$

Proof Setting $\hat{u} := Eg$ and $\hat{v} := E\tau_{\gamma}v$ as well as using (2.4) proves the inequalities. We note that (2.3) follows directly from the corollary as well.

In the subsequent sections we introduce and discuss some different functionals Φ and corresponding constants c_{Φ} .

2.1 First estimate

For any $y \in \mathsf{D}(\Omega)$ and any $w \in \overset{\circ}{\mathsf{H}}_{-1}^{1}(\Omega)$ we have

$$\langle \operatorname{div} y, w \rangle_{\Omega} + \langle y, \nabla w \rangle_{\Omega} = 0.$$
 (2.5)

Combining (2.1) and (2.5) we obtain for all $w \in \overset{\circ}{\mathsf{H}}_{-1}^1(\Omega)$ and all $y \in \mathsf{D}(\Omega)$

$$\langle A\nabla(u-v), \nabla w \rangle_{\Omega} = \langle f + \operatorname{div} y, w \rangle_{\Omega} + \langle y - A\nabla v, \nabla w \rangle_{\Omega} =: \Phi(w). \tag{2.6}$$

By Cauchy-Schwarz' inequality, (1.3) with $c_N := 2/(N-2)$ and (1.5) we estimate the right hand side $\Phi(w)$ of (2.6) as follows:

$$\begin{split} |\langle f + \operatorname{div} y, w \rangle_{\Omega}| &\leq \|f + \operatorname{div} y\|_{1,\Omega} \|w\|_{-1,\Omega} \leq c_{N} \|f + \operatorname{div} y\|_{1,\Omega} \|\nabla w\|_{\Omega} \\ &\leq \frac{c_{N}}{\sqrt{c_{A}}} \|f + \operatorname{div} y\|_{1,\Omega} \|\nabla w\|_{A,\Omega} \\ |\langle y - A\nabla v, \nabla w \rangle_{\Omega}| &\leq \|y - A\nabla v\|_{A^{-1},\Omega} \|\nabla w\|_{A,\Omega} \end{split} \tag{2.8}$$

By Corollary 4 we arrive at the following result.

Proposition 5 Let u, v be as in Theorem 3. Then

$$\|\nabla(u-v)\|_{A,\Omega} \le \frac{c_N}{\sqrt{c_A}} \|f + \operatorname{div} y\|_{1,\Omega} + \|y - A\nabla v\|_{A^{-1},\Omega} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)}, \qquad (2.9)$$

where y is an arbitrary vector field in $D(\Omega)$.

Remark 6 If v satisfies the prescribed boundary condition, then (2.9) implies

$$\|\nabla(u-v)\|_{A,\Omega} \le \frac{c_N}{\sqrt{c_A}} \|f + \operatorname{div} y\|_{1,\Omega} + \|y - A\nabla v\|_{A^{-1},\Omega}. \tag{2.10}$$

The estimates (2.9) and (2.10)) show that deviations from exact solutions of exterior boundary value problems have the same structure as for problems in bounded domains, namely they contain weighted residuals of basic relations with weights given by constants in the corresponding embedding inequalities.

2.2 Second estimate

Assume that Ω is decomposed into two subdomains Ω_i and Ω_e with interface $\Gamma := \partial \Omega_e$ (see Figure 1) and that the fields $y \in \mathsf{D}(\Omega)$ exactly satisfy the relation

$$\operatorname{div} y + f = 0 \quad \text{in } \Omega_e. \tag{2.11}$$

In particular, this situation may arise if the source term f has compact support and g is represented (in the exterior domain Ω_e) as a linear combination of solenoidal fields having proper decay at infinity. In this case, the estimate of Proposition 5 turns trivially to

$$\|\nabla(u-v)\|_{A,\Omega} \le c_o \|f + \operatorname{div} y\|_{\Omega_i} + \|y - A\nabla v\|_{A^{-1},\Omega} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)}, \tag{2.12}$$

which holds for all $y \in D(\Omega)$ additionally satisfying (2.11), where the weight constant is

$$c_o := \frac{c_N(1 + ||r||_{\infty,\Omega_i})}{\sqrt{c_A}},\tag{2.13}$$

which follows directly from

$$||f + \operatorname{div} y||_{1,\Omega} = ||f + \operatorname{div} y||_{1,\Omega_i} \le |\rho|_{\infty,\Omega_i} ||f + \operatorname{div} y||_{\Omega_i} \le (1 + |r|_{\infty,\Omega_i}) ||f + \operatorname{div} y||_{\Omega_i}.$$

But we also may derive another estimate. We rewrite (2.7) and use Cauchy-Schwarz' inequality in Ω_i

$$\left| \langle f + \operatorname{div} y, w \rangle_{\Omega} \right| = \left| \langle f + \operatorname{div} y, w \rangle_{\Omega_i} \right| \le \| f + \operatorname{div} y \|_{\Omega_i} \| w \|_{\Omega_i}$$
 (2.14)

and estimate

$$\|w\|_{\Omega_i} \le c_{\Omega_i} \|\nabla w\|_{\Omega_i} \le \frac{c_{\Omega_i}}{\sqrt{c_A}} \|\nabla w\|_{A,\Omega}.$$
 (2.15)

Here c_{Ω_i} denotes a Poincare/Friedrich constant associated with the bounded domain Ω_i , i.e. the best constant of the inequality

$$\|\varphi\|_{\Omega_i} \leq c_{\Omega_i} \, \|\nabla\varphi\|_{\Omega_i} \quad \forall \, \varphi \in \left\{ \psi \in \mathsf{H}^1(\Omega_i) \mid \tau_{\partial \, \Omega_i} \psi|_{\gamma} = 0 \text{ on } \gamma \right\},$$

where $\tau_{\partial\Omega_i}: \mathsf{H}^1(\Omega_i) \to \mathsf{H}^{1/2}(\partial\Omega_i)$ denotes the trace operator. In this case, we have again (2.12) but now with the (optional) weight constant

$$c_o := \frac{c_{\Omega_i}}{\sqrt{c_A}}. (2.16)$$

We note that the constant (2.13) may also be achieved by (2.7) and the argument (2.14) if we replace the estimate (2.15) by

$$\|w\|_{\Omega_i} \le (1+|r|_{\infty,\Omega_i}) \|w\|_{-1,\Omega_i} \le (1+|r|_{\infty,\Omega_i}) \|w\|_{-1,\Omega} \le \frac{c_N}{\sqrt{c_A}} (1+|r|_{\infty,\Omega_i}) \|\nabla w\|_{A,\Omega}.$$

We summarize and get our second a posteriori error estimate.

Proposition 7 For all $y \in D(\Omega)$ with (2.11) we have

$$\|\nabla(u-v)\|_{A,\Omega} \le c_o \|f + \operatorname{div} y\|_{\Omega_i} + \|y - A\nabla v\|_{A^{-1},\Omega} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)},$$

where c_o is defined either by (2.13) or by (2.16).

Remark 8 In general, the number c_{Ω_i} will be smaller and thus provides a better bound than $c_N(1 + \|r\|_{\infty,\Omega_i})$. On the other hand, the number $c_N(1 + \|r\|_{\infty,\Omega_i})/\sqrt{c_A}$ is an easily computable upper bound for the best possible constant c_o .

2.3 Third estimate

Let y_i and y_e be the restrictions of some $y \in L^2(\Omega)$ to Ω_i and Ω_e , respectively. Assuming $y_i \in D(\Omega_i)$ and $y_e \in D(\Omega_e)$ but not necessarily $y \in D(\Omega)$ we use the equations

$$\langle y_i, \nabla w \rangle_{\Omega_i} + \langle \operatorname{div} y_i, w \rangle_{\Omega_i} = \langle \tau_{n,\Gamma} y_i, \tau_{\Gamma} w \rangle_{\Gamma},$$
 (2.17)

$$\langle y_e, \nabla w \rangle_{\Omega_e} + \langle \operatorname{div} y_e, w \rangle_{\Omega_e} = -\langle \tau_{n,\Gamma} y_e, \tau_{\Gamma} w \rangle_{\Gamma},$$
 (2.18)

which hold for all $w \in \overset{\circ}{\mathsf{H}}^{1}_{-1}(\Omega)$ and in the sense of the traces $\tau_{\Gamma} : \mathsf{H}^{1}_{-1}(\Omega) \to \mathsf{H}^{1/2}(\Gamma)$ and $\tau_{n,\Gamma} : \mathsf{D}(\Omega_{i}) \to \mathsf{H}^{-1/2}(\Gamma)$ respectively $\tau_{n,\Gamma} : \mathsf{D}(\Omega_{e}) \to \mathsf{H}^{-1/2}(\Gamma)$. At this point we assume that the interface Γ is Lipschitz (in order to guarantee that the traces are well defined). By $\langle \varphi, \psi \rangle_{\Gamma}$ we denote the duality product of $\mathsf{H}^{-1/2}(\Gamma)$ and $\mathsf{H}^{1/2}(\Gamma)$. We recall that the normal traces $\tau_{n,\Gamma}y_{i}$ and $\tau_{n,\Gamma}y_{e}$ possess weak surface divergences in $\mathsf{H}^{-1/2}(\Gamma)$ as well. If $y \in \mathsf{D}(\Omega)$, then div $y_{i} = \mathrm{div}\,y$ in Ω_{i} and div $y_{e} = \mathrm{div}\,y$ in Ω_{e} . Hence, in this case adding (2.17) and (2.18) we obtain by (2.5)

$$\langle \tau_{n,\Gamma} y_i - \tau_{n,\Gamma} y_e, \tau_{\Gamma} w \rangle_{\Gamma} = \langle y, \nabla w \rangle_{\Omega_i} + \langle \operatorname{div} y, w \rangle_{\Omega} = 0$$

for all $w \in \overset{\circ}{\mathsf{H}}_{-1}^1(\Omega)$. Therefore, we get

$$\tau_{n,\Gamma} y_i = \tau_{n,\Gamma} y_e$$

for all $y \in \mathsf{D}(\Omega)$ since τ_{Γ} is surjective.

On our way to find Φ like in (2.6) we now insert (2.17), (2.18) instead of (2.5) into (2.1) and obtain

$$\langle A\nabla(u-v), \nabla w \rangle_{\Omega} = \langle f + \operatorname{div} y_i, w \rangle_{\Omega_i} + \langle f + \operatorname{div} y_e, w \rangle_{\Omega_e}$$

$$+ \langle y - A\nabla v, \nabla w \rangle_{\Omega} + \langle \tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i, \tau_{\Gamma} w \rangle_{\Gamma} =: \Phi(w).$$
(2.19)

The third term of $\Phi(w)$ will be estimated by (2.8) and for the last term we may use the continuity of the trace operator τ_{Γ} in combination with a Poincare/Friedrich estimate, i.e.

$$\|\tau_{\Gamma}\varphi\|_{\mathsf{H}^{1/2}(\Gamma)} \le c_{\Gamma} \|\nabla\varphi\|_{A,\Omega} \quad \forall \varphi \in \overset{\circ}{\mathsf{H}}_{-1}^{1}(\Omega), \tag{2.20}$$

and obtain

$$\left| \left\langle \tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i, \tau_{\gamma} w \right\rangle_{\Gamma} \right| \leq \left\| \tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i \right\|_{\mathsf{H}^{-1/2}(\Gamma)} \left\| \tau_{\Gamma} w \right\|_{\mathsf{H}^{1/2}(\Gamma)}$$

$$\leq c_{\Gamma} \left\| \tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i \right\|_{\mathsf{H}^{-1/2}(\Gamma)} \left\| \nabla w \right\|_{A,\Omega}.$$

$$(2.21)$$

To estimate the second term of $\Phi(w)$ we again use (1.3) and (1.5) and obtain

$$\left| \langle f + \operatorname{div} y_e, w \rangle_{\Omega_e} \right| \leq \| f + \operatorname{div} y_e \|_{1,\Omega_e} \| w \|_{-1,\Omega_e} \leq \| f + \operatorname{div} y_e \|_{1,\Omega_e} \| w \|_{-1,\Omega}
\leq \frac{c_N}{\sqrt{c_A}} \| f + \operatorname{div} y_e \|_{1,\Omega_e} \| \nabla w \|_{A,\Omega}.$$
(2.22)

Considering the first (and last) term of $\Phi(w)$ we have once more at least two options as in section 2.2 to obtain the estimate

$$\left| \langle f + \operatorname{div} y_i, w \rangle_{\Omega_i} \right| \le c_o \| f + \operatorname{div} y_i \|_{\Omega_i} \| \nabla w \|_{A,\Omega} \tag{2.23}$$

with c_o defined either by (2.13) or (2.16).

Finally with (2.19) and (2.8), (2.21), (2.22), (2.23) we get by Corollary 4 the third estimate.

Proposition 9 For all $y \in L^2(\Omega)$ with $y_i \in D(\Omega_i)$ and $y_e \in D(\Omega_e)$ we have

$$\|\nabla(u-v)\|_{A,\Omega} \le c_o \|f + \operatorname{div} y_i\|_{\Omega_i} + \frac{c_N}{\sqrt{c_A}} \|f + \operatorname{div} y_e\|_{1,\Omega_e} + \|y - A\nabla v\|_{A^{-1},\Omega}$$

$$+ c_\Gamma \|\tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i\|_{\mathsf{H}^{-1/2}(\Gamma)} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)}$$

$$(2.24)$$

with c_o from Proposition 7. The right hand side of (2.24) vanishes if and only if v coincides with u and y with $A\nabla u$.

Remark 10 There are many ways to deduce (2.20). We just mention that $\tau_{\Gamma}\varphi$ can be considered as a trace of a function defined in Ω_i or Ω_e or even of a function, which is just defined in a small neighborhood of Γ . Thus, we may adjust the constant c_{Γ} according to our needs.

Remark 11 This estimate suggests even a solution method: We construct approximations using locally supported trial functions in Ω_i , e.g. FEM, and utilize global approximations properly behaving at infinity for Ω_e . These two types of approximations are usually difficult to meet together exactly on the artificial boundary Γ . However, Proposition 9 shows that this is not required because we can use instead the penalty term with known penalty factor c_{Γ} . In addition, we have one more parameter, the 'radius' of the interface Γ . Since Γ is artificial and arbitrary we can use this parameter in the algorithm in order to obtain better results.

Remark 12 At this point we shall note that all our estimates are sharp, which easily can be seen by setting $v := u \in H^1_{-1}(\Omega)$ and $y := A\nabla u \in D(\Omega)$.

Remark 13 In Propositions 5, 7, 9 we can always replace the last summand of the right hand side by $2 \|\nabla(\hat{u} - \hat{v})\|_{A,\Omega}$ or $2 \|\nabla E(g - \tau_{\gamma}v)\|_{A,\Omega}$ using Theorem 3 and Corollary 4.

3 Upper bounds in dimension N=2

Of course, Theorem 3 holds for N=2 as well and the modifications on the estimates depend just on the Poincare/Friedrich estimate and thus they are obvious using the proper Cauchy-Schwarz inequality. We achieve

Proposition 14 Let $\Omega \subset \mathbb{R}^2$, such that $\mathbb{R}^2 \setminus \Omega$ contains the unit ball.

(i) For all $y \in D(\Omega)$

$$\|\nabla(u-v)\|_{A,\Omega} \le \frac{2}{\sqrt{c_A}} \|r \ln r(f+\operatorname{div} y)\|_{\Omega} + \|y-A\nabla v\|_{A^{-1},\Omega} + 2c_{\gamma} \|g-\tau_{\gamma} v\|_{\mathsf{H}^{1/2}(\gamma)}.$$

(ii) For all $y \in D(\Omega)$ with div y + f = 0 in Ω_e

$$\|\nabla(u-v)\|_{A,\Omega} \le c_o \|f + \operatorname{div} y\|_{\Omega_i} + \|y - A\nabla v\|_{A^{-1},\Omega} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)},$$
where $c_o = \min \left\{ 2 \|r \ln r\|_{\infty,\Omega_i}, c_{\Omega_i} \right\} / \sqrt{c_A}.$

(iii) For all $y \in L^2(\Omega)$ with $y_i \in D(\Omega_i)$ and $y_e \in D(\Omega_e)$

$$\|\nabla(u-v)\|_{A,\Omega} \le c_o \|f + \operatorname{div} y_i\|_{\Omega_i} + \frac{2}{\sqrt{c_A}} \|r \ln r(f + \operatorname{div} y_e)\|_{1,\Omega_e} + \|y - A\nabla v\|_{A^{-1},\Omega} + c_\Gamma \|\tau_{n,\Gamma} y_e - \tau_{n,\Gamma} y_i\|_{\mathsf{H}^{-1/2}(\Gamma)} + 2c_\gamma \|g - \tau_\gamma v\|_{\mathsf{H}^{1/2}(\gamma)}.$$

Analogously, Remarks 8, 10, 11, 12, 13 hold.

A Appendix

A.1 Lower bounds for the error

We note by a standard variational argument

$$\|\nabla(u-v)\|_{A,\Omega}^2 = \sup_{y \in \mathsf{L}^2(\Omega)} \left(2 \left\langle A \nabla(u-v), y \right\rangle_{\Omega} - \|y\|_{A,\Omega}^2 \right).$$

Thus, we obtain for all $w \in \mathsf{H}^1_{-1}(\Omega)$ the estimate

$$\begin{split} \|\nabla(u-v)\|_{A,\Omega}^2 &\geq 2 \left\langle A\nabla(u-v), \nabla w \right\rangle_{\Omega} - \|\nabla w\|_{A,\Omega}^2 \\ &= 2 \left\langle A\nabla u, \nabla w \right\rangle_{\Omega} - \left\langle A\nabla(2v+w), \nabla w \right\rangle_{\Omega}, \end{split}$$

which is sharp since one can put w = u - v. But to exclude the unknown exact solution u from the right hand side we need $w \in \overset{\circ}{\mathsf{H}}^1_{-1}(\Omega)$ since then by (1.4)

$$\|\nabla(u-v)\|_{A,\Omega}^2 \ge 2\langle f, w \rangle_{\Omega} - \langle A\nabla(2v+w), \nabla w \rangle_{\Omega}. \tag{A.1}$$

But this estimate is no longer sharp because we can not put w = u - v anymore. In fact, with $A\nabla u \in \mathsf{D}(\Omega)$ and div $A\nabla u = -f$ we get for $w \in \mathsf{H}^1_{-1}(\Omega)$

$$\langle A\nabla u, \nabla w \rangle_{\Omega} = \langle f, w \rangle_{\Omega} + \langle \tau_{n,\gamma} A\nabla u, \tau_{\gamma} w \rangle_{\gamma}.$$

Hence, we obtain the estimate

$$\left\|\nabla(u-v)\right\|_{A,\Omega}^{2} \geq 2\left\langle f,w\right\rangle_{\Omega} - \left\langle A\nabla(2v+w),\nabla w\right\rangle_{\Omega} + 2\left\langle \tau_{n,\gamma}A\nabla u,\tau_{\gamma}w\right\rangle_{\gamma}$$

for all $w \in \mathsf{H}^1_{-1}(\Omega)$, which is sharp and coincides with (A.1) if $w \in \mathsf{H}^1_{-1}(\Omega)$. But the unknown exact solution u still appears on the right-hand side, i.e. the normal trace of $A\nabla u$ on γ . Furthermore, if $\langle \tau_{n,\gamma} A\nabla u, \tau_{\gamma} w \rangle_{\gamma} > 0$ then (A.1) can not be sharp.

A.2 Poincare type estimates for exterior domains

We introduce the radial derivative $\partial_r := \xi \cdot \nabla$, where $\xi(x) := x/r(x)$. Furthermore, B_{ε} and S_{ε} denote the open ball and sphere of radius ε centered at the origin in \mathbb{R}^N , respectively. We will use the ideas of [4, Lemma 4.1] and [1, Poincare's estimate III, p. 57] with some minor useful modifications.

Lemma 15 Let $\Omega \subset \mathbb{R}^N$, $N \geq 1$, be a domain and $\beta \in \mathbb{R}$. For all $u \in \overset{\circ}{\mathsf{C}}^{\infty}(\Omega)$ the following Poincare estimates hold:

(i) If $\beta > 1 - N/2$ then

$$\left(2\beta+N-2\right)\left\Vert r^{\beta-1}u\right\Vert _{\Omega}\leq2\left\Vert r^{\beta}\,\partial_{r}\,u\right\Vert _{\Omega}.$$

(ii) Let $B_1 \subset \mathbb{R}^N \setminus \Omega$. If $\beta \geq (3-N)/2$ or $\beta \leq 1-N/2$ then

$$|2\beta + N - 3| \left\| \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega} \le 2 \left\| r^{\beta} \partial_r u \right\|_{\Omega}.$$

(iii) If N=1 then

$$|2\beta - 1| \|(1+r)^{\beta-1}u\|_{\Omega} \le 2 \|(1+r)^{\beta} \partial_r u\|_{\Omega} + |2\min\{0, 2\beta - 1\}|^{1/2} |u(0)|,$$

where u will be extended by zero to \mathbb{R} .

For the estimates derived in this paper it suffices to set $\beta = 0$. In this particular case, the above lemma implies

Corollary 16 Let $\Omega \subset \mathbb{R}^N$, $N \geq 1$, be a domain. For all $u \in \overset{\circ}{\mathsf{C}}^{\infty}(\Omega)$ the following Poincare estimates hold:

(i) If $N \geq 3$ then

$$\|u\|_{-1,\Omega} \le \|u/(1+r)\|_{\Omega} \le \|u/r\|_{\Omega} \le \frac{2}{N-2} \|\partial_r u\|_{\Omega} \le \frac{2}{N-2} \|\nabla u\|_{\Omega}.$$

(ii) If N = 2 and $B_1 \subset \mathbb{R}^2 \setminus \Omega$ then

$$||u/(r \ln r)||_{\Omega} \le 2 ||\partial_r u||_{\Omega} \le 2 ||\nabla u||_{\Omega}$$

(iii) If N = 1 then

$$\|u\|_{-1,\Omega} \le \|u/(1+r)\|_{\Omega} \le 2 \|\partial_r u\|_{\Omega} + \sqrt{2}|u(0)| \le 2 \|u'\|_{\Omega} + \sqrt{2}|u(0)|.$$

Hence, if $\Omega \subset \mathbb{R}_{\pm}$ we have

$$||u||_{-1,\Omega} \le ||u/(1+r)||_{\Omega} \le 2 ||\partial_r u||_{\Omega} \le 2 ||u'||_{\Omega}.$$

Remark 17 Of course, by continuity all these estimates extend to appropriate weighted H¹-Sobolev spaces.

Proof Let $\Omega \subset \mathbb{R}^N$, $N \geq 1$, be a domain and $u \in \overset{\circ}{\mathsf{C}}^{\infty}(\Omega)$. By partial integration we get for all $\alpha \in \mathbb{R}$ and $\varepsilon > 0$

$$2\int_{\Omega\setminus B_{\varepsilon}} r^{\alpha} u \, \partial_{r} u \, d\lambda = \int_{\Omega\setminus B_{\varepsilon}} r^{\alpha} \, \partial_{r} |u|^{2} \, d\lambda$$
$$= -(\alpha + N - 1) \int_{\Omega\setminus B_{\varepsilon}} r^{\alpha - 1} |u|^{2} \, d\lambda - \varepsilon^{\alpha} \int_{S_{\varepsilon}} |u|^{2} \, d\sigma.$$

Thus, for all $\gamma \in \mathbb{R}$ and $\beta := (\alpha + 1)/2$

$$\begin{split} \left\| r^{\beta} \, \partial_{r} \, u + \gamma r^{\beta - 1} u \right\|_{\Omega \backslash B_{\varepsilon}}^{2} \\ &= \left\| r^{\beta} \, \partial_{r} \, u \right\|_{\Omega \backslash B_{\varepsilon}}^{2} + |\gamma|^{2} \left\| r^{\beta - 1} u \right\|_{\Omega \backslash B_{\varepsilon}}^{2} + 2 \gamma \underbrace{\left\langle r^{\beta} \, \partial_{r} \, u, r^{\beta - 1} u \right\rangle_{\Omega \backslash B_{\varepsilon}}}_{= \int_{\Omega \backslash B_{\varepsilon}} r^{\alpha} u \, \partial_{r} \, u \, \mathrm{d}\lambda} \\ &= \left\| r^{\beta} \, \partial_{r} \, u \right\|_{\Omega \backslash B_{\varepsilon}}^{2} + \gamma (\gamma - 2\beta - N + 2) \left\| r^{\beta - 1} u \right\|_{\Omega \backslash B_{\varepsilon}}^{2} - \gamma \varepsilon^{2\beta - 1} \int_{S_{\varepsilon}} |u|^{2} \, \mathrm{d}\sigma. \end{split}$$

Now the left hand side of this equality converges by the monotone convergence theorem. Since $r^{\nu} \in \mathsf{L}^1(U_1)$, if and only if $\nu > -N$, and $|\int_{S_{\varepsilon}} |u|^2 \, \mathrm{d}\sigma| \leq c\varepsilon^{N-1}$ the right hand side converges for $\beta > 1 - N/2$ by Lebesgue's dominated convergence theorem in \mathbb{R} . Hence, for $\varepsilon \to 0$ we obtain

$$\|r^{\beta} \partial_r u + \gamma r^{\beta-1} u\|_{\Omega}^2 = \|r^{\beta} \partial_r u\|_{\Omega}^2 + \gamma (\gamma - 2\beta - N + 2) \|r^{\beta-1} u\|_{\Omega}^2.$$

Choosing $\gamma := 2\beta + N - 2 > 0$ we finally get by the triangle inequality

$$\gamma \| r^{\beta - 1} u \|_{\Omega} \le 2 \| r^{\beta} \partial_r u \|_{\Omega}.$$

Since we are especially interested in the case $\beta = 0$ this estimate is only applicable in dimensions $N \geq 3$.

For N=1 we proceed as follows: For all $\alpha \in \mathbb{R}$ we have

$$2\int_{\mathbb{R}_{\pm}} (1+r)^{\alpha} u \, \partial_r \, u \, d\lambda = \pm 2\int_{\mathbb{R}_{\pm}} (1\pm t)^{\alpha} u(t) u(t)' \, dt$$
$$= \pm 2\int_{\mathbb{R}_{+}} (1\pm t)^{\alpha} \left(|u(t)|^2 \right)' \, dt = -\alpha \int_{\mathbb{R}_{+}} (1\pm t)^{\alpha-1} |u(t)|^2 \, dt - |u(0)|^2$$

and thus

$$2 \int_{\mathbb{R}} (1+r)^{\alpha} u \, \partial_r \, u \, d\lambda = -\alpha \int_{\mathbb{R}} (1+r)^{\alpha-1} |u(t)|^2 \, d\lambda - 2|u(0)|^2.$$

Hence, for all $\gamma \in \mathbb{R}$ and $\beta := (\alpha + 1)/2$

$$\begin{aligned} & \left\| (1+r)^{\beta} \, \partial_{r} \, u + \gamma (1+r)^{\beta-1} u \right\|_{\Omega}^{2} \\ &= \left\| (1+r)^{\beta} \, \partial_{r} \, u \right\|_{\Omega}^{2} + |\gamma|^{2} \, \left\| (1+r)^{\beta-1} u \right\|_{\Omega}^{2} + 2\gamma \underbrace{\left\langle (1+r)^{\beta} \, \partial_{r} \, u, (1+r)^{\beta-1} u \right\rangle_{\Omega}}_{= \int_{\Omega} (1+r)^{\alpha} u \, \partial_{r} \, u \, d\lambda} \\ &= \left\| (1+r)^{\beta} \, \partial_{r} \, u \right\|_{\Omega}^{2} + \gamma (\gamma - 2\beta + 1) \, \left\| (1+r)^{\beta-1} u \right\|_{\Omega}^{2} - 2\gamma |u(0)|^{2}. \end{aligned}$$

As before the triangle inequality and the choice $\gamma := 2\beta - 1$, but now without any restrictions on β , lead to

$$\begin{split} |\gamma| \, \big\| (1+r)^{\beta-1} u \big\|_{\Omega} & \leq \big\| (1+r)^{\beta} \, \partial_{r} \, u \big\|_{\Omega} + \Big(\big\| (1+r)^{\beta} \, \partial_{r} \, u \big\|_{\Omega}^{2} - 2\gamma |u(0)|^{2} \Big)^{1/2} \\ & \leq 2 \, \big\| (1+r)^{\beta} \, \partial_{r} \, u \big\|_{\Omega} + |2 \min\{0,\gamma\}|^{1/2} \, |u(0)|. \end{split}$$

The remaining case N=2 requires the use of logarithms. Moreover, the origin is now a problematic singularity, which has to be removed from our domain. Therefore, we may assume $B_1 \subset \mathbb{R}^N \setminus \Omega$ and $N \geq 1$, having N=2 in mind. We start once more for all $\alpha \in \mathbb{R}$ with

$$2\int_{\Omega} \frac{r^{\alpha}}{\ln r} u \, \partial_r u \, d\lambda = \int_{\Omega} \frac{r^{\alpha}}{\ln r} \, \partial_r |u|^2 \, d\lambda$$
$$= -(\alpha + N - 1) \int_{\Omega} \frac{r^{\alpha - 1}}{\ln r} |u|^2 \, d\lambda + \int_{\Omega} \frac{r^{\alpha - 1}}{\ln^2 r} |u|^2 \, d\lambda.$$

Now our usual procedure gives for $\gamma \in \mathbb{R}$ and $\beta := (\alpha + 1)/2 \ge 0$

$$\begin{split} \left\| r^{\beta} \, \partial_r \, u + \gamma \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2 \\ &= \left\| r^{\beta} \, \partial_r \, u \right\|_{\Omega}^2 + |\gamma|^2 \left\| \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2 + 2\gamma \underbrace{\left\langle r^{\beta} \, \partial_r \, u, \frac{r^{\beta - 1}}{\ln r} u \right\rangle_{\Omega}}_{= \int_{\Omega} \frac{r^{\alpha}}{\ln r} u \, \partial_r \, u \, \mathrm{d}\lambda \\ &= \left\| r^{\beta} \, \partial_r \, u \right\|_{\Omega}^2 + \gamma (\gamma + 1) \left\| \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2 - \gamma (N + 2\beta - 2) \left\| \frac{r^{\beta - 1}}{\sqrt{\ln r}} u \right\|_{\Omega}^2. \end{split}$$

Thus, for $\gamma(N+2\beta-2) \geq 0$ we can estimate

$$\left\| r^{\beta} \partial_r u + \gamma \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2 \le \left\| r^{\beta} \partial_r u \right\|_{\Omega}^2 + \gamma (\gamma - 2\beta - N + 3) \left\| \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2,$$

which leads to the estimate

$$\left\| r^{\beta} \, \partial_r \, u + \gamma \frac{r^{\beta - 1}}{\ln r} u \right\|_{\Omega}^2 \le \left\| r^{\beta} \, \partial_r \, u \right\|_{\Omega}^2$$

if we set $\gamma := 2\beta + N - 3$ with the additional constraint $\gamma(\gamma + 1) \ge 0$, i.e. $\gamma \ge 0$ or $\gamma \le -1$. Finally, again by the triangle inequality

$$|\gamma| \left\| \frac{r^{\beta-1}}{\ln r} u \right\|_{\Omega} \le 2 \left\| r^{\beta} \partial_r u \right\|_{\Omega}$$

follows for all $\beta \geq (3-N)/2$ or $\beta \leq (2-N)/2$.

Acknowledgements The authors express their gratitude to the Department of Mathematical Information Technology of the University of Jyväskylä (Finland) for financial support.

References

- [1] Leis, R., Initial Boundary Value Problems in Mathematical Physics, Teubner, Stuttgart, (1986).
- [2] Repin, S., 'A posteriori error estimates for variational problems with uniformly convex functionals', *Math. Comp.*, 69, (230), (2000), 481-500.

- [3] Repin, S., A posteriori estimates for partial differential equations, Radon Series Comp. Appl. Math., ISBN: 978-3-11-019153-0, Walter de Gruyter, Berlin, (2008).
- [4] Saranen, J., Witsch, K.-J., 'Exterior boundary value problems for elliptic equations', Ann. Acad. Sci. Fenn. Math., 8, (1), (1983), 3-42.

Dirk Pauly

University of Duisburg-Essen Faculty for Mathematics Campus Essen

Universitätsstr. 2 45117 Essen Germany

e-mail: dirk.pauly@uni-due.de

and

University of Jyväskylä Faculty of Information Technology Department of

Mathematical Information Technology

P.O. Box 35 (Agora) FI-40014 Jyväskylä

Finland

e-mail: dirk.pauly@jyu.fi

Sergey Repin

V.A. Steklov Mathematical Institute

St. Petersburg Branch

Fontanka 27

191011 St. Petersburg

Russia

e-mail: repin@pdmi.ras.ru

University of Jyväskylä

Faculty of Information Technology

Department of

Mathematical Information Technology

P.O. Box 35 (Agora) FI-40014 Jyväskylä

Finland

e-mail: serepin@jyu.fi